



Northland Power Inc. on behalf of Northland Power Solar Abitibi L.P. Toronto, Ontario


Natural Heritage Evaluation of Significance Report

Abitibi Solar Project

H334844-0000-07-124-0244 Rev. 1 October 18, 2012

Disclaimer

This report has been prepared by or on behalf of Northland Power Inc. for submission to the Ontario Ministry of the Environment as part of the Renewable Energy Approval process. The content of this report is not intended for the use of, nor is it intended to be relied upon by, any other person. Neither Northland Power Inc. nor any of its directors, officers, employees, agents or consultants has any liability whatsoever for any loss, damage or injury suffered by any third party arising out of, or in connection with, their use of this report.





Project Report

October 18, 2012

Northland Power Inc. Abitibi Solar Project

Natural Heritage Evaluation of Significance Report Table of Contents

| 1. | Introduction | 3 |
|----|--|----|
| | 1.1 Project Description | 3 |
| | 1.2 Legislative Requirements | 3 |
| | 1.2.1 Records Review Report | |
| | 1.2.2 Site Investigation Report | |
| | 1.2.3 Evaluation of Significance Report | |
| | 1.3 Input to Evaluation of Significance from Consultation Activities | 11 |
| | 1.3.1 Public Consultation | |
| | 1.3.2 Aboriginal Consultation | |
| | 1.3.3 Municipal/Local Authority Consultation | 11 |
| | 1.4 Evaluation of Significance Report Format | 11 |
| 2. | Summary of Results of Records Review and Site Investigation | 12 |
| 3. | Wildlife Habitat | 12 |
| | 3.1 Evaluation Criteria and Guidelines for Wildlife Habitat, and Determination of Significance | 14 |
| | 3.1.1 Solar Panel Project Location | |
| | 3.1.1.1 Seasonal Concentration Areas | |
| | 3.1.1.2 Specialized Wildlife Habitat | |
| | 3.1.1.3 Habitat for Species of Conservation Concern | |
| | 3.1.1.4 Animal Movement Corridors | 21 |
| | 3.1.2 Transmission Line Project Location | 22 |
| | 3.2 Date of Beginning and Completion of Evaluation | 23 |
| | 3.3 Overall Conclusion | 23 |
| | 3.4 Name and Qualifications of Evaluator | |
| 4. | Wetlands | 25 |
| | 4.1 Solar Panel Project Location | 25 |
| | 4.2 Transmission Line Project Location | |
| 5. | Conclusions | 31 |
| 6. | References | 32 |
| Αp | pendix A Natural Resource Solutions Inc., Wetland Evaluations | |



List of Tables

| Table 2.1 | Natural Features on and within 120 m of the Project Location | 12 |
|------------|--|----|
| Table 4.1 | Wetland Characteristics and Ecological Functions of Wetlands within 120 m of the | |
| | Transmission Line Project Location | 27 |
| Table 5.1 | Significant Natural Features on and within 120 m of the Project Location | 31 |
| | List of Figures | |
| Figure 1.1 | Project Location and Significant Natural Heritage Features | 5 |
| Figure 1.2 | Transmission Line Project Location (Eastern Half) – | |
| | Generalized Candidate Significant Natural Heritage Features | 7 |
| Figure 1.3 | Transmission Line Project Location (Western Half) – | |
| | Generalized Candidate Significant Natural Heritage Features | 9 |
| Figure 3.1 | Solar Panel Project Location and Evaluation of Significance – Survey Locations | 18 |





1. Introduction

1.1 Project Description

Northland Power Solar Abitibi L.P. (hereinafter referred to as "Northland") is proposing to develop a Class 3 10-megawatt (MW) ground mounted solar photovoltaic (Solar PV) facility in the District of Cochrane. This Project, known as the Abitibi Solar Project, is hereafter referred to as "Abitibi" or the "Project."

The Project location is comprised of two primary components. The first part of the Project is the location of the solar panels, including access roads, inverters, transformers, fencing, etc, and is hereafter referred to as the "solar panel Project location" The solar panel Project location is approximately 98 hectares (ha) in size and located on Lots 14 and 15, Concession 8 of the Town of Cochrane. The solar panel Project location is situated on Glackmeyer Concession Road 9 (shown in Figure 1.1).

The second part of the Project is the approximately 20 km transmission line from the solar panel Project location to the connection point west of the Project location near Hunta, ON, as well as associated transition structure and switching station. This portion of the project is referred to as the transmission line Project location, with locations shown in Figures 1.2 and 1.3.

1.2 Legislative Requirements

Ontario Regulation (O. Reg.) 359/09 – Renewable Energy Approvals Under Part V.O.1 of the Act, made under the Environmental Protection Act identifies the Renewable Energy Approval (REA) requirements for renewable energy projects in Ontario. Ground-mounted solar facilities with a name plate capacity greater than 10 kilowatts (kW) are classified as Class 3 solar facilities and require a REA in accordance with Section 4 of O. Reg. 359/09.

Section 24(1) of O. Reg. 359/09 requires proponents of Class 3 solar projects to undertake a natural heritage assessment consisting of a records review report, site investigation report and an evaluation of significance report for each natural feature identified during the records review and site investigation.

Natural Features are defined in Section 1(1) of O. Reg. 359/09 to be all or part of

- a) an area of natural and scientific interest (ANSI) (earth science)
- b) an ANSI (life science)
- c) a coastal wetland
- d) a northern wetland
- e) a southern wetland
- f) a valleyland
- g) a wildlife habitat, or
- h) a woodland.







In respect of woodlands and valleylands, Section 1(1) of O. Reg. 359/09 requires that these features be located south and east of the Canadian Shield as shown in Figure 1 in the Provincial Policy Statement issued under Section 3 of the *Planning Act*. This figure shows that the proposed Project location is located on the Canadian Shield, and therefore valleylands and woodlands as defined by O. Reg. 359/09 cannot be located on the Project location.

1.2.1 Records Review Report

Section 25 of the REA Regulation requires proponents of Class 3 solar projects to undertake a natural heritage records review to identify "whether the project is

- (a) in a natural feature
- (b) within 50 m of an area of natural and scientific interest (earth science)
- (c) within 120 m of a natural feature that is not an area of natural or scientific interest (earth science)." (O. Reg. 359/09, s. 25, Table).

Subsection 2 of Section 30 of the REA Regulation requires the proponent to prepare a report "setting out a summary of the records searched and the results of the analysis" (O. Reg. 359/09). The Natural Heritage Records Review Report (Hatch Ltd., 2012a) was prepared to meet these requirements.

1.2.2 Site Investigation Report

Section 26 of the REA Regulation requires proponents of Class 3 solar projects to undertake a natural heritage site investigation for the purpose of determining

- whether the results of the analysis summarized in the (natural heritage records review) report prepared under Subsection 25(3) are correct or require correction, and identifying any required corrections
- whether any additional natural features exist, other than those that were identified in the (natural heritage records review) report prepared under Subsection 30(2)
- the boundaries, located within 120 m of the Project location, of any natural feature that was identified in the records review or the site investigation
- the distance from the Project location to the boundaries determined under clause (c).


The Natural Heritage Site Investigation Report (Hatch Ltd., 2012b) was prepared to meet these requirements.

1.2.3 Evaluation of Significance Report

Section 27 of the REA Regulation requires proponents of Class 3 solar projects to undertake an evaluation of significance (EOS) for natural heritage features identified during the records review and site investigation and prepare a report that sets out

- a determination of whether the natural feature is
 - provincially significant
 - significant








Blank back








back







back





- not significant
- not provincially significant
- a summary of the evaluation criteria or procedures used to make the determinations
- the name and qualifications of any person who applied to evaluation criteria or procedures.

This EOS Report for the natural features identified within 120 m of the Project has been prepared to meet these requirements.

1.3 Input to Evaluation of Significance from Consultation Activities

As required by Section 27 of O.Reg. 359/09, the evaluation of significance must consider information obtained through consultation with the public, aboriginal communities and municipalities and local authorities. Results of these consultation activities in relation to the evaluation of significance are discussed below.

1.3.1 Public Consultation

A public meeting has been held in association with this Project; notices for the meeting were published in the local newspaper. In addition, landowners within 120 m of the Project location were mailed notices of the proposed Project and meeting dates.

To date, no information relating to natural features relevant to the evaluation of significance has been obtained through these consultation activities.

1.3.2 Aboriginal Consultation

Aboriginal communities identified by the Ministry of the Environment as communities to be consulted through the Renewable Energy Approval process have been mailed letters requesting information relating to the Project, along with meeting notices and copies of the Project Description Report.

To date, no information relating to natural features relevant to the evaluation of significance has been obtained through these consultation activities.

1.3.3 Municipal/Local Authority Consultation

Meetings have been held with staff of the Town of Cochrane and Hunta Local Roads Board. In addition, the Town and Roads Board has received notices of the public meetings and copies of the Project Description Report.

To date, no information relating to natural features relevant to the evaluation of significance has been obtained through these consultation activities.

1.4 Evaluation of Significance Report Format

Section 1 of this EOS has identified the legislative requirements for an EOS under the REA Regulation and identified the reasons why an EOS is required for the Project. Section 2 provides a summary of the results of the records review and site investigation. Section 3 provides the EOS for wildlife habitat, and Section 4 provides the EOS for the wetland. Section 5 identifies the conclusions of the EOS, and the references are provided in Section 6.





2. Summary of Results of Records Review and Site Investigation

As stated above, natural features requiring an evaluation of significance are identified through the records review (Hatch Ltd., 2011a) and site investigation (Hatch Ltd., 2011b) required under Sections 25 and 26 of the REA Regulation, respectively. These studies have already been completed, and the results are summarized in Table 2.1. This Report provides the evaluations for the features identified in Table 2.1.

Table 2.1 Natural Features on and within 120 m of the Project Location


| Natural Feature | al Feature Project Location | |
|------------------------------|-----------------------------|-----|
| Solar Panel Project Location | on | |
| ANSI – Earth Science | No | No |
| ANSI – Life Science | No | No |
| Wetland | Yes | Yes |
| Wildlife Habitat | Yes | Yes |
| Transmission Line Project | Location | |
| ANSI – Earth Science | No | No |
| ANSI – Life Science | No | No |
| Wetland | No | Yes |
| Wildlife Habitat | No | Yes |

3. Wildlife Habitat

Several types of candidate significant wildlife habitats were identified during the site investigation:

- Solar Panel Project Location
 - Waterfowl Nesting habitat
 - Habitat for area-sensitive species
 - Wetlands supporting amphibian breeding habitat
 - Habitat for species of conservation concern, including
 - Common Nighthawk Habitat
 - Olive-sided Flycatcher Habitat
 - Canada Warbler Habitat
 - Vaccinium ovalifoliuym habitat
 - Carex wiegandii habitat
 - Carex haydenii habitat
 - Animal movement corridor
- Transmission Line Project Location
 - Generalized Characterized Candidate Significant Wildlife Habitat







- Seasonal Concentration Areas
 - o Winter deer yards/moose late winter habitat
 - o Waterfowl stopover and staging areas
 - o Waterfowl nesting sites
- Specialized Wildlife Habitats
 - o Area-sensitive woodland/shrubland/grassland habitats
 - o Moose aquatic feeding areas
 - o Old growth or mature forest stands
 - o Woodlands supporting amphibian breeding habitat
 - o Wetlands supporting amphibian breeding habitat
 - o Mink, otter, marten and fisher denning sites
 - o Specialized raptor nesting habitat
 - o Seeps and springs
- Habitat for Species of Conservation Concern
 - o Northern Long-eared Bat
 - o Red-necked Grebe
 - o Short-eared Owl
 - o Common Nighthawk
 - o Canada Warbler
 - o Bald Eagle
 - o Olive-Sided Flycatcher
 - o Vaccinium ovalifolium
 - o Scirpus heterochaetus
 - o Carex wiegandii
 - o Carex tetanica
 - o Carex Ioliacea
 - o Carex haydenii
- Animal Movement Corridors associated with several waterbodies within 120 m of the Project location







3.1 Evaluation Criteria and Guidelines for Wildlife Habitat, and Determination of Significance

The criteria processes outlined in the Ministry of Natural Resources (MNR) Natural Heritage Assessment Guide (NHAG) (MNR, 2011) and Significant Wildlife Habitat Technical Guide (SWHTG) (MNR, 2000) are used to evaluate the significance of wildlife habitat. The specific criteria used in the evaluation from these sources are discussed by habitat type below.

3.1.1 Solar Panel Project Location

3.1.1.1 Seasonal Concentration Areas

3.1.1.1.1 Waterfowl Nesting Habitat

In accordance with Appendix D of the NHAG, waterfowl nesting habitat identified within 120 m of the Solar Panel Project location during the site investigations is considered to be "Generalized Candidate Significant Wildlife Habitat". As such this feature will be carried forward to the Environmental Impact Study.

3.1.1.2 Specialized Wildlife Habitat

Criteria for evaluation of specialized habitat for wildlife are identified within Table Q-2 of Appendix Q of the SWTHG. The criteria that were considered during the evaluation of these features are discussed in respect of the individual features below.

3.1.1.2.1 Wetlands supporting amphibian breeding habitat

Wetlands supporting amphibian breeding habitat were identified within the wetland communities around the creek within 120 m west of the Project location. In order to evaluate the significance of wetlands supporting amphibian breeding habitat, amphibian calling surveys were completed at a point within the wetland community on two separate occasions. Surveys were completed in accordance with the protocols outlined in the Marsh Monitoring Program, which consists of 180 deg, 3-minute point counts, completed either after sunset or after 2200 hours. Survey locations are shown in Figure 3.1. Details of the surveys are provided below:

- Site Investigation 1
 - Date, Times and Duration of Site Investigation

Date: May 18, 2011

Start Time: 2115

■ End Time: 2235

Duration: 1 hour 20 minutes

Weather Conditions During Site Investigation

■ Temperature: 7-10°C

Cloud Cover: Clear sky

Name and Qualifications of Person Conducting Site Investigation







- This site investigation was completed by Levi Snook and Shelley Potter:
 - Levi Snook is an Environmental Scientist with experience conducting
 environmental assessments on proposed hydroelectric, wind, and solar energy
 sites. He has diplomas in environmental science from Sir Sandford Fleming
 College and a degree in biology from Trent University. He has expertise in
 terrestrial assessments in support of Natural Heritage studies that include
 conducting Ecological Land Classifications, as well as wildlife inventories,
 including amphibian and reptile surveys.
 - Shelley Potter is an environmental professional with a marine and freshwater biology honours graduate from the University of Guelph. Previous work and internships have provided experience in the fields of environmental science, sustainable development, water conservation and analysis, fresh water biology, marine mammal biology, Ichthyology and Oceanography. Shelley recently completed an internship with the University of Queensland working with Dr. Mike Noad at the Humpback Whale Acoustic Research Collaboration. Marine Mammal Observing experience, acoustic recording experience and ability to geographically track migration patterns of humpback whales using a theodolite and Cyclops computer program was acquired. Shelley has also recently participated in terrestrial and aquatic field surveys for various renewable energy projects in Ontario.
- Site Investigation 2
 - Date, Times and Duration of Site Investigation

Date: June 21, 2011

Start Time: 2000

■ End Time: 2200

Duration: 2 hours

Weather Conditions During Site Investigation

Temperature: 15 °C

Beaufort Wind: 1

Cloud Cover: 5%

- Name and Qualifications of Person Conducting Site Investigation
 - Names and qualifications of NRSI staff conducting the site investigations are provided in Appendix B.

During the first site investigation, American Toad, Wood Frogs and Spring Peeper were heard within the wetland, while only spring peepers were recorded during the second site investigation.







The results of these site investigations were then used to assess the criteria for significant wetlands supporting amphibian breeding habitat:

- Provision of significant wildlife habitat The wetland community is also considered to be a
 candidate significant animal movement corridor and waterfowl nesting habitat, and therefore this
 criteria is met.
- Degree of permanence It is expected that water is permanently found within the creek, therefore this criteria is met.
- Species diversity of pond Three species of frog (Spring Peeper, American Toad, Wood Frog)
 were recorded during amphibian surveys. Therefore, species diversity of the ponds is considered
 to be moderate.
- Presence of rare species No rare species were identified during the baseline surveys.
- Size and number of ponds The wetland community is relatively large and therefore this criteria is met.
- Diversity of submergent and emergent vegetation A diversity of submergent and emergent vegetation was not recorded from the wetland community.
- Presence of shrubs, logs at edge of pond Both tall and low shrubs were recorded within the
 wetland community, therefore this criteria is met.
- Adjacent forest habitat Portions of the wetland community occur adjacent to forest communities, therefore this criteria is met.
- Water quality Water quality is unknown.
- Level of disturbance Active agricultural operations occur on either side of the wetland community.


Therefore, as the criteria for provision of significant wildlife habitat, degree of permanence, species diversity, size, presence of shrubs and adjacent forest habitat have been met, this feature is determined to be a significant wetland supporting amphibian breeding habitat.

3.1.1.2.2 Habitat for Area-Sensitive Shrubland Species

Area-sensitive shrubland birds were assessed through a random area search of suitable habitats during the breeding season. The search area is shown in Figure 3.1. Details of this survey are provided below (note: duration includes area searches of all habitat types).

- Date, Times and Duration of Site Investigation
 - Date: June 21, 2011
 - Start Time: 0530
 - End Time: 0800
 - Duration: 2.5 hours
- Weather Conditions During Site Investigation







• Temperature: 13°C

Beaufort Wind: 0 to 2

Cloud Cover: 90%

Name and Qualifications of Person Conducting Site Investigation

 Names and qualifications of NRSI staff conducting the site investigations are provided in Appendix A.

Of the birds detected, none are considered to be area-sensitive shrubland species. Therefore, this habitat is not considered to be significant habitat for area-sensitive species, and further comparison to the criteria is not required.

3.1.1.2.3 Habitat for Area-Sensitive Woodland Species

Area-sensitive woodland birds were assessed through a random area search of suitable habitats during the breeding season. The search area is shown in Figure 3.1. Details of this survey are provided in Section 3.1.1.2.2

Of the birds species recorded, several are considered to be area-sensitive woodland species; Black-and White Warbler, Ovenbird, Black-throated Green Warbler and American Redstart. Singing male Black-and-white Warbler, Ovenbird and Black-throated Green Warbler were recorded within the upland forest community in the central portion of the Project location. Singing male American Redstart were recorded within the woodland communities on and within 120 m of the southern portions of the Project location.

These results were then compared against the criteria for area-sensitive species:

- Presence of rare, uncommon or declining species None of these species are a rare, uncommon
 or declining species, and therefore this criteria is not met.
 - Overall area of site –The upland forest community within the central portion of the Project location is 15 ha in size, and therefore this criteria is not met. The forest community on and within 120 m of the southern portion of the Project site is part of a larger network of forest communities, and therefore this criteria is met.
- Area of forest interior contained within the forest stand With respect to the upland forest community within the central portion of the Project location there is no interior forest habitat, and therefore this criteria is not met. With respect to the woodlands on and within 120 m of the southern portion of the Project location, there is more than 10 ha of interior forest found present within the stand, and therefore this criteria is met.
- Age and tree composition of forest stand The upland forest community within the central
 portion of the Project location does not contain an abundance of large, mature trees. With
 respect to the woodlands on and within 120 m of the southern portion of the Project location, an
 array of tree composition was noted within the stand, with most trees mid-aged. Therefore, this
 criteria is met.







- Amount of vertical stratification of site –Vertical stratification was not noted within the forest communities, and therefore this criteria is not met.
- Amount of contiguous closed-canopy/open areas in forest stand The forest communities have a
 relatively closed canopy, and therefore this criteria is met.
- Degree of disturbance on site Given the adjacent agricultural activity, this criteria is not met.
- Amount of adjacent residential development There are occasional residences, but no true residential development, therefore this criteria is met.
- Current representation of habitat in planning area This habitat is abundantly available within the planning area, therefore this criteria is not met.
- Provision of significant wildlife habitat There are no other candidate significant wildlife
 habitats identified in association with these communities, therefore this criteria is not met.

Therefore, as the woodland in the central portion of the Project location did not meet either the criteria for size or interior forest, this habitat is not considered to be significant.

With respect to the woodlands on and within 120 m of the southern portion of the Project location, though several of the criteria have been met, as American Redstart are an abundant species both within the region and the province, and since breeding habitat for American Redstart is also abundantly available across the province, areas of habitat on and within 120 m of the Project location are not considered to be significant wildlife habitat

3.1.1.3 Habitat for Species of Conservation Concern

3.1.1.3.1 Canada Warbler

Area searches of woodland habitats, as previously described in Section 3.1.1.2.3 did not result in any observations of Canada Warbler. As a result, it is determined that they are not present on or within 120 m of the Project location.

3.1.1.3.2 Olive-sided Flycatcher

Area searches of shrubland and woodland habitats, as previously described in Sections 3.1.1.2.2 and 2.1.1.2.3, respectively, did not result in any observations of Olive-sided Flycatcher. As a result, it is determined that they are not present on or within 120 m of the Project location.


3.1.1.3.3 Common Nighthawk

Evening bird surveys were completed in conjunction with the second site investigation for wetlands supporting amphibian breeding habitat (see Section 3.1.1.2.1 for details of timing and weather conditions). Survey locations are shown in Figure 3.1. No Common Nighthawk were recorded during the surveys on or within 120 m of the Project location.

3.1.1.3.4 Carex haydenii

This species was not detected during vegetation surveys of suitable habitats on and within 120 m of the Project location. Details of vegetation surveys have been previously identified in the Natural Heritage Site Investigations Report (Hatch, 2012b).







back of Figure 3.1







Carex wiegandii

This species was not detected during vegetation surveys of suitable habitats on and within 120 m of the Project location. Details of vegetation surveys have been previously identified in the Natural Heritage Site Investigations Report (Hatch, 2012b).

3.1.1.3.5 Vaccinium ovalifolium

This species was not detected during vegetation surveys of suitable habitats on and within 120 m of the Project location. Details of vegetation surveys have been previously identified in the Natural Heritage Site Investigations Report (Hatch, 2012b).

3.1.1.4 Animal Movement Corridors

A candidate significant animal movement was identified in association with the creek and the associated riparian habitat on and within 120 m of the Project location. Evaluation of animal movement corridors is identified within Section 8.7 of the SWHTG. The criteria for significance are outlined in Table Q-4 of Appendix Q in the SWHTG, and are provided below along with the evaluation for these features:

- Importance of areas to be linked by corridor The corridor links Lauzon Lake with waterbodies
 further north, likely providing linkage between breeding and foraging areas for a variety of
 wildlife species, therefore this criteria is met.
- Dimensions of corridor The corridor near the Project location varies in width from 250 to 300 m wide, and therefore this criteria is met.
- Continuity of corridor The corridor is broken by a road, and therefore this criteria is not met.
- Habitat and habitat structure of corridor The corridor consists of a range of habitats from marshland, to thicket swamp, to coniferous swamp, to upland forest, this criteria is met.
- Species found in corridor or presumed to be using corridor It is assumed that a diverse array of species would use the corridor and therefore this criteria is met.
- Risk of mortality for species using corridor There is a moderate risk of mortality for species
 using the corridor given the presence of a road crossing, though not well travelled, and open
 agricultural lands adjacent to the corridor providing for ease of predator movement. Therefore,
 this criteria is not met.
- Opportunity for protection As this feature is associated with a watercourse, opportunity for protection is good and therefore this criteria is met.
- Provision of other related values (such as erosion protection) As this corridor includes riparian
 habitats, it provides protection for soil erosion and water quality, as well as for foraging
 opportunities for other wildlife species. Therefore, this criteria is met.

Therefore, as several criteria have been met, the corridor is determined to be a significant animal movement corridor.







3.1.2 Transmission Line Project Location

In accordance with Appendix D of the NHAG, all wildlife habitat identified within 120 m of the transmission line Project location during the site investigations are considered to be "Generalized Candidate Significant Wildlife Habitat". As such the features listed below will be carried forward to the Environmental Impact Study:

- Seasonal Concentration Areas
 - Winter deer yards/moose late winter habitat
 - Waterfowl stopover and staging areas
 - Waterfowl nesting sites
- Specialized Wildlife Habitats
 - Area-sensitive woodland/shrubland/grassland habitats
 - Moose aquatic feeding areas
 - Old growth or mature forest stands
 - Woodlands supporting amphibian breeding habitat
 - Wetlands supporting amphibian breeding habitat
 - Mink, otter, marten and fisher denning sites
 - Specialized raptor nesting habitat
 - Seeps and springs
- Habitat for Species of Conservation Concern
 - Northern Long-eared Bat
 - Red-necked Grebe
 - Short-eared Owl
 - Common Nighthawk
 - Canada Warbler
 - Bald Eagle
 - Olive-Sided Flycatcher
 - Vaccinium ovalifolium
 - Scirpus heterochaetus
 - Carex wiegandii
 - Carex tetanica
 - Carex Ioliacea







- Carex haydenii
- Animal Movement Corridors associated with several waterbodies within 120 m of the Project location

3.2 Date of Beginning and Completion of Evaluation

The evaluation of wildlife habitat commenced with records review in May 2010 and is finalized with the completion of this Report in February 2012. Site investigations were completed in association with this evaluation on August 24, 2010, and May 18, June 23, and June 24, 2011.

3.3 Overall Conclusion

Based on the evaluation above, the following significant wildlife habitat features were identified:

- Solar panel Project location
 - Seasonal Concentration Areas
 - Waterfowl Nesting Habitat (Generalized Characterized Candidate Significant Wildlife Habitat)
 - Specialized Wildlife Habitats
 - Wetlands supporting amphibian breeding habitat
 - Animal movement corridor associated with the creek and riparian habitat
- Transmission line Project location
 - Generalized Characterized Candidate Significant Wildlife Habitat
 - Seasonal Concentration Areas
 - o Winter deer yards/moose late winter habitat
 - o Waterfowl stopover and staging areas
 - Waterfowl nesting sites
 - Specialized Wildlife Habitats
 - o Area-sensitive woodland/shrubland/grassland habitats
 - o Moose aquatic feeding areas
 - o Old growth or mature forest stands
 - o Woodlands supporting amphibian breeding habitat
 - o Wetlands supporting amphibian breeding habitat
 - o Mink, otter, marten and fisher denning sites
 - Specialized raptor nesting habitat
 - o Seeps and springs







- Habitat for Species of Conservation Concern
 - o Northern Long-eared Bat
 - Red-necked Grebe
 - Short-eared Owl
 - o Common Nighthawk
 - o Canada Warbler
 - o Bald Eagle
 - o Olive-Sided Flycatcher
 - Vaccinium ovalifolium
 - o Scirpus heterochaetus
 - o Carex wiegandii
 - o Carex tetanica
 - o Carex Ioliacea
 - o Carex haydenii
- Animal Movement Corridors associated with several waterbodies within 120 m of the Project location

3.4 Name and Qualifications of Evaluator

Evaluations of wildlife habitat were completed by Sean K. Male of Hatch.

Sean K. Male, M.Sc. is a Terrestrial Ecologist specializing in assessments of terrestrial habitat, flora and fauna. Sean received his Bachelors of Science (Honours) in Biology from Queen's University, where he completed his Honour's thesis under Dr. Raleigh J. Robertson, studying the impacts of nestbox density in Tree Swallows (*Tachycineta bicolor*) on nest-building behaviour. He then completed a Master's of Science degree in the Watershed Ecosystem Graduate Program at Trent University under Dr. Erica Nol. Sean's thesis focussed on examining the impacts of a Canadian diamond mine on a population of breeding passerines. For his thesis, Sean spent two summers in the Canadian arctic studying populations of Lapland Longspurs (*Calcarius lapponicus*) around the Ekati Diamond Mine, located 300 km northeast of Yellowknife. While at Trent, Sean participated in the Northern Saw-whet Owl (*Aegoius acadicus*) Migration Banding Project at the Oliver Centre. Following his time at Trent, Sean participated in the Landscape Monitoring Program, participating in a study of the impacts of woodlot size on breeding birds.

Sean joined Hatch as a Terrestrial Ecologist in 2006. Since joining Hatch, Sean has participated in several environmental assessments, REAs and other regulatory approvals for hydro, wind and solar power developments as the terrestrial biologist specializing in field investigations identifying flora and fauna species, including species of significance. He has developed and implemented baseline monitoring and impact assessment programs for both terrestrial wildlife and plant communities,





including detailed bird and bat studies for several wind power developments, including the proposed 100-MW Coldwell wind power development near Marathon, Ontario, a proposed 20-MW facility near Port Dover, Ontario, and a proposed 110-MW wind facility in southwestern Ontario. Sean has also conducted terrestrial and wetland vegetation surveys for several proposed hydropower projects totalling over 40 MW in southern and northern Ontario and has participated in fisheries surveys for several of these projects.

4. Wetlands

4.1 Solar Panel Project Location

The evaluation of the wetland communities was completed separately and can be found in Appendix A. The conclusion of the wetland evaluation was that these communities were provincially significant.

4.2 Transmission Line Project Location

There are no wetlands identified on the Transmission Line Project Location; however, there are a number of wetlands identified within 120 m of this Project location. These wetlands have been identified to be associated with 10 wetland complexes.

Two of these wetland complexes have been previously assessed as a provincially significant wetland (see Section 4.1 and Hatch, 2012c). In accordance with Appendix C of the Natural Heritage Assessment Guide, the eight remaining wetland complexes are assumed to be provincially significant wetlands. The characteristics of these eight wetland communities that must be documented as per the requirements of Appendix C are identified in Table 4.1.









 Table 4.1
 Wetland Characteristics and Ecological Functions of Wetlands within 120 m of the Transmission Line Project Location

| Ecological Function | Wetland Catch Basin 2 | Wetland Catch Basin 3 | Wetland Catch Basin 4 | Wetland Catch Basin 5 | Wetland Catch Basin 6 | Wetland Catch Basin 7 | Wetland Catch Basin 8 |
|--|---|---|---|---|---|---|--|
| Drainage Basin Size (ha) | 2015 | 1432 | 807 | | 1061 | 1496 | 1424 |
| Actual Wetland Size (ha) (within 120 m of transmission line) | 47.67 | 29.47 | 34.57 | 1.6 | 30.1 | 33.1 | 22.23 |
| Wetland Type | Within 120 m – Marsh Swamp/thicket swamp communities also present within catch basin | Within 120 m – Swamp Marsh/thicket swamp communities also present within catch basin | Within 120 m – Swamp Marsh/thicket swamp communities also present within catch basin | Within 120 m – Marsh Swamp/thicket swamp communities also present within catch basin | Within 120 m – Swamp Marsh/thicket swamp communities also present within catch basin | Within 120 m – Swamp Marsh/thicket swamp communities also present within catch basin | Within 120 m – Marsh Swamp/thicket swamp communities also present within catch basin |
| Site Type | Palustrine/Riverine | Palustrine/Riverine | Palustrine/Riverine | Palustrine/Riverine | Palustrine/Riverine | Palustrine/Riverine | Palustrine/Riverine/ Lacustrine |
| Vegetation Communities (within 120 m of transmission line) | gcM; cS; tsS | cS; gcM | cS; tsS | gcM | cS; tsS; gcM | cS; tsS; gcM | gcM; tsS; cS |
| Proximity to other wetlands | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km | Hydrologically connected by surface water to other wetlands (different dominant wetland type) or open lake or river within 1.5 km |
| Interspersion | Medium to High | Medium to High |
| Open Water Types | Type 3: Open water occupies 5-25% of the wetland area, occurring in ponds of various sizes; vegetation occurs in dense patches or diffuse open stands. | Type 1: Open water occupies < 5% of the wetland area. | Type 2: Open water occupies 5 to 25% of the wetland area, occurring in a central area | Type 2: Open water occupies 5-25% of the wetland area, occurring in a central area | Type 3: Open water occupies 5-25% of the wetland area, occurring in ponds of various sizes; vegetation occurs in dense patches or diffuse open stands | Type 3: Open water occupies 5-25% of the wetland area, occurring in ponds of various sizes; vegetation occurs in dense patches or diffuse open stands | Type 5: Open water occupies 26-75% of the wetland area, occurring in a pattern where small ponds and "embayments" are common |
| Flood Attenuation (total) | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. | None of the wetland communities located within 120 m of the Project location are considered to be isolated communities, however isolated wetland communities may be found elsewhere within the catchment basin, more than 120 m from the Project location. Wetland vegetation communities along watercourses within 120 m of the Project would provide stormwater retention, and therefore flood attenuation benefits. |







| Ecological Function | Wetland Catch Basin 2 | Wetland Catch Basin 3 | Wetland Catch Basin 4 | Wetland Catch Basin 5 | Wetland Catch Basin 6 | Wetland Catch Basin 7 | Wetland Catch Basin 8 |
|--------------------------------------|---|---|--|--|--|---|---|
| Water Quality Improvement (total) | Given the presence of: • tertiary roadways and corridors within the catchment basin • agricultural and forestry activities within the upland areas the wetland communities are determined to provide moderate water quality improvement functions | Given the presence of: a secondary roadway within the catchment basin tertiary roadways and corridors within the catchment basin agricultural and forestry activities within the upland areas a small pit within the catchment basin the wetland communities are determined to provide moderate water quality improvement functions | Given the presence of: • tertiary roadways and corridors within the catchment basin • agricultural and forestry activities within the upland areas the wetland communities are determined to provide low water quality improvement functions | Given the presence of: • tertiary roadways and corridors within the catchment basin • agricultural and forestry activities within the upland areas the wetland communities are determined to provide low water quality improvement functions | Given the presence of: a major hydro corridor from the Long Sault hydroelectric station within the catchment basin tertiary roadways and corridors within the catchment basin agricultural and forestry activities within the upland areas the wetland communities are determined to provide moderate to high water quality improvement functions | Given the presence of: a major hydro corridor from the Long Sault hydroelectric station within the catchment basin tertiary roadways and corridors within the catchment basin forestry activities within the upland areas the wetland communities are determined to provide moderate to high water quality improvement functions | Given the presence of: a major hydro corridor from the Long Sault hydroelectric station within the catchment basin tertiary roadways and corridors within the catchment basin agricultural and forestry activities within the upland areas the wetland communities are determined to provide moderate to high water quality improvement functions |
| Shoreline Erosion Control | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine wetland communities within (and beyond) 120 m of the Project location provide high shoreline erosion control functions | Thicket shrub and meadow marsh communities associated with the palustrine/riverine and lacustrine wetland communities found within the catchment basin provide high shoreline erosion control functions |
| Groundwater Recharge (Total) | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. | Wetlands within 120 m of the Project location are primarily palustrine/riverine, and therefore provide limited to medium potential for groundwater recharge, primarily during annual flooding events, such as spring freshet. The lacustrine wetland communities around the south shore of Lower Deception Lake would provide no groundwater recharge potential. |
| Species Rarity (Total) | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species = none | No rare species noted during 2011 surveys within the wetland. • Breeding Habitat for Endangered or Threatened Species |







| Ecological Function | Wetland Catch Basin 2 | Wetland Catch Basin 3 | Wetland Catch Basin 4 | Wetland Catch Basin 5 | Wetland Catch Basin 6 | Wetland Catch Basin 7 | Wetland Catch Basin 8 |
|--|--|--|--|--|--|--|--|
| | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species of Special Status = none | = none Traditional Migration or Feeding Areas for an Endangered or Threatened Species = none Provincially Significant Plant and Animal Species = none Regionally Significant Species = none Locally Significant Species = none Species = none Species of Special Status = none |
| Significant Features and Habitats (Total) | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none | Colonial Waterbirds = none Winter Cover for Wildlife = none Waterfowl Staging and/or Molting Area = none Waterfowl Breeding = none |
| Fish Habitat (Total) | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/movement function as fish travel to and from various habitat areas. | No specific fish community or habitat assessment work has been conducted. The watercourses and associated riparian wetlands within the community are deemed to provide spawning, nursery and residence habitat for the fish community, as well as some migration/ movement function as fish travel to and from various habitat areas. |





Blank back





5. Conclusions

Results of the EOS are summarized in Table 5.1. Based on the EOS outlined above, there are significant natural features present on and within 120 m of the Project location. The locations of these features are shown in Figures 1.1 to 1.3.

An environmental impact study conducted according to the requirements of Section 38(2) of O. Reg. 359/09 will be required in order to construct the Project within 120 m of these significant natural features.

Table 5.1 Significant Natural Features on and within 120 m of the Project Location

| N | latural Feature | Project Location | Adjacent Lands (within 120 m) | |
|-----------------------------|--------------------------|------------------|---|--|
| Solar Par | nel Project Location | | | |
| SIGNIFICANT | Wildlife Habitat | Yes | Yes | |
| | Wetland | Yes | Yes | |
| N I | Earth Science ANSI | No | No | |
| PROVINCIALLY SIGNIFICANT | Life Science ANSI | No | No | |
| Transmis | sion Line Project Locati | on | | |
| SIGNIFICANT | Wildlife Habitat | No | Yes (generalized candidate significant wildlife habitat) | |
| ALLY | Wetland | No | Yes (2 evaluated, 8 assumed provincially significant) | |
| \(\zeta \) | Earth Science ANSI | No | No | |
| PROVINCIALLY SIGNIFICANT | Life Science ANSI | No | No | |







6. References

Hatch Ltd. 2012a. Abitibi Solar Project – Natural Heritage Records Review Report. Prepared for Northland Power Inc. on behalf of Northland Power Solar Abitibi L.P.

Hatch Ltd. 2012b. Abitibi Solar Project – Natural Heritage Site Investigations Report. Prepared for Northland Power Inc. on behalf of Northland Power Solar Abitibi L.P.

Hatch Ltd. 2012c. Long Lake Solar Project – Natural Heritage Site Investigations Report. Prepared for Northland Power Inc. on behalf of Northland Power Solar Long Lake L.P.

Ministry of Natural Resources (MNR). 2011. Natural Heritage Assessment Guide for Renewable Energy Projects. Toronto: Queen's Printer for Ontario. 248 pp.

MNR. 2000. Significant Wildlife Habitat Technical Guide. 151p.





Appendix A

Natural Resource Solutions Inc. (NRSI)
Wetland Evaluation



1247B

February 22, 2012

Mr. Sean Male Hatch Energy 4342 Queen Street, Suite 500 Niagara Falls, ON L2E 7J7

Dear Mr. Male,

RE: Abitibi and Martin's Meadow Solar Project

Summary of Wetland & Upland Vegetation Mapping,

Breeding Bird and Amphibian Call Surveys

Summary of Surveys

On behalf of Natural Resource Solutions Inc., I am pleased to provide the following which documents the work completed at the above noted solar project being proposed by Northland Power.

The objectives of this assignment were to complete vegetation mapping, amphibian surveys, breeding bird, and evening bird surveys.

Appendix I includes a list of study team members and their roles.

Vegetation

On site vegetation mapping occurred on June 22, 2011 (0900 – 1600hrs, weather 15℃, sunny, 0% cloud cover, wind – Beaufort scale 1). The standard Ontario Wetland Evaluation System (OWES) (OMNR 1993) was used by a Certified Wetland Evaluator to map and describe on-site wetlands, as well as wetlands within 120m of the project site.

In addition, a catchment basin boundary was identified that included the on-site wetlands. All wetlands in the catchment basin were also mapped and described using OWES June 21 to June 24, 2011. In this case, land access and the extent of the lands required that the mapping be completed using aerial photography supplemented with field checks of wetland polygons at strategic locations (primarily roadside).

The standard Ecological Land Classification (ELC) (Lee et al. 1998; Lee 2008) was also used by a Certified ELC staff to describe polygons outside of OWES and Forest Ecosystem Classification (FEC) (Taylor et al. 2000).

Please see Appendix II for a list of polygon labels.

The wetlands within the catchment basin were evaluated using the standard OWES system for northern Ontario. A copy of the completed evaluation, including mapping, is included in Appendix III.

Amphibian Call Monitoring

On site amphibian call surveys were completed on June 21, 2011 (2000-2200hrs, weather 15° C, 5% cloud cover, wind – Beaufort scale 3 to 4, water temperature 19° C). The standard Marsh Monitoring Protocol (Bird Studies Canada 2009) was used in which 3 minute point counts at predetermined stations.

At the Abitibi site nothing was heard at Station 1, which was determined to be marginal amphibian habitat since no water or frog habitat was present. Three spring peepers (*Pseudacris crucifer crucifer*) were heard northeast of Station 1B (approximately 100m distance). Two spring peepers were heard north of Station 2 (approximately 100m distance). At the Martin's Meadow site, nothing was heard at Station 2. No standing water or frog habitat is present. A second station was chosen, Station 3, to replace monitoring at Station 2 which was at a sedge marsh with pockets of standing water. No amphibians were heard.

The field data forms are included in Appendix IV.

Breeding Bird Surveys

On site breeding bird surveys were completed June 21, 2011 (0530 – 0800hrs, weather 13°C, 90% cloud cover, wind – Beaufort scale 0 to 2) using the standard Ontario Breeding Bird methodology (Cadman et al. 2007).

The following species were observed during that period:

| Species Observed | Observed | Possible | Probable | Confirmed |
|---|----------|----------|----------|-----------|
| Alder Flycatcher (Empidonax alnorum) | | S | | |
| American Crow (Corvus brachyrhynchos) | Х | | | |
| American Goldfinch (Carduelis tristis) | | Н | | |
| American Redstart (Setophaga ruticilla) | | S | | |
| American Robin (Turdus migratorius) | | S | | |
| Black-and-white Warbler (Mniotilta varia) | | S | | |
| Black-throated Green Warbler (Dendroica virens) | | S | | |
| Blue Jay (Cyanocitta cristata) | | S | | |
| Common Loon (Gavia immer) | Х | | | |
| Hermit Thrush (Catharus guttatus) | | S | | |
| Ovenbird (Seiurus aurocapillus) | | S | | |
| Red-eyed Vireo (Vireo olivaceus) | | S | | |
| Sandhill Crane (Grus canadensis) | | | | FY |
| Song Sparrow (Melospiza melodia) | | S | | |
| Tennessee Warbler (Vermivora peregrine) | | S | | |
| White-throated Sparrow (Zonotrichia albicollis) | | S | | |
| Yellow Warbler (Dendroica petechia) | | S | | |
| Yellow-rumped Warbler (Dendroica cronoata) | Х | | | |

The following species were observed within the Martin's Meadow area:

| Species Observed | Observed | Possible | Probable | Confirmed |
|---|----------|----------|----------|-----------|
| American Crow (Corvus brachyrhynchos) | Х | | | |
| American Goldfinch (Carduelis tristis) | | S | | |
| American Redstart (Setophaga ruticilla) | | S | | |
| American Robin (Turdus migratorius) | | S | | |
| Black-throated Green Warbler (Dendroica virens) | Х | | | |
| Northern Cardinal (Cardinalis cardinalis) | | S | | |
| Ovenbird (Seiurus aurocapillus) | | S | | |
| Red-eyed Vireo (Vireo olivaceus) | | S | | |
| Sandhill Crane (Grus canadensis) | | S | | |
| Savannah Sparrow (Passerculus sandwichensis) | | S | | |
| Veery (Catharus fuscescens) | | S | | |
| White-throated Sparrow (Zonotrichia albicollis) | | S | | |

Observed

X Species observed in its breeding season with no evidence of breeding

Possible

- H Species observed in its breeding season in suitable nesting habitat
- S Singing male present of breeding calls heard in breeding season in suitable nesting habitat Probable
- P Pair observed in their breeding season in suitable nesting habitat
- T Permanent territory presumed through registration of territorial song on at least 2 days, one week or more apart at the same place
- D Courtship or display between a male and female or 2 males including courtship feeding and copulation
- V Visiting probable nest site
- A Agitated behaviour or anxiety calls of an adult
- B Brood patch on adult female or cloacal protuberance on adult male
- N Nest building or excavation of nest site

Confirmed

- DD Distraction display or injury feigning
- NU Used nest or egg shell found (occupied/laid this season)
- FY Recently fledged young or downy young
- AE Adults leaving or entering nest site in circumstances indicating occupied nest
- FS Adult carrying faecal sac
- CF Adult carrying food for young
- NE Nest containing eggs
- NY Nest with young seen or heard

Other species observed on site included:

Red Fox (Vulpes vulpes)

Evening Bird Surveys

Surveys for birds that are primarily active in the evening were conducted on June 21, 2011 (2000-2200hrs, weather 15%, 5% cloud cover, wind – Beaufort scale 3 to 4). The survey followed standard monitoring protocols developed for species such as whippoor-will and common nighthawk (the two focus species for this survey) (OMNR 2011). No nighthawks (Chordeiles sp.) or whip-poor-wills (*Caprimulgus vociferous*) were observed during evening surveys.

Other species observed during evening surveys included:

American Robin (*Turdus migratorius*)
Hermit Thrush (*Catharus guttatus*)
Sandhill Crane (*Grus canadensis*)
Veery (*Catharus fuscescens*)
White-throated Sparrow (*Zonotrichia albicollis*)

Red Fox (Vulpes vulpes)

References

- Bird Studies Canada. 2009. Marsh Monitoring Program Participant's Handbook for Surveying Amphibians. 2009 Edition. Published by Bird Studies Canada in Cooperation with Environment Canada and the U.S. Environmental Protection Agency. February 2009.
- Cadman, M.D., D.A. Sutherland, G.G. Beck, D. Lepage and A.R. Couturier. 2007. Atlas of the Breeding Birds of Ontario. Available online at: http://www.birdsontario.org/atlas/index.jsp
- Lee, H.T., W.D. Bakowsky, J. Riley, J. Bowles, M. Puddister, P. Uhlig and S. McMurray. 1998. Ecological Land Classification for Southern Ontario: First Approximation and its Application. Ontario Ministry of Natural Resources, Southcentral Science Section, Science Development and Transfer Branch. SCSS Field Guide FG-02.
- Lee, H. 2008. Southern Ontario Ecological Land Classification Vegetation Type List. Ontario Ministry of Natural Resources: London, Ontario.
- Ontario Ministry of Natural Resources. 1993. Ontario Wetland Evaluation System. Northern Manual. Revised 1994 & 2002)
- Ontario Ministry of Natural Resources. 2011. Northeast Nightjar Survey Techniques (Draft). Sudbury District MNR
- Taylor, K.C. et al. 2000. A Field Guide to Forest Ecosystems of Northeastern Ontario. 2nd Edition. NEST Field Guide FG-001.

Appendix I Team Members

Appendix I

| Team Member | Qualification | Role |
|------------------|-----------------------------|-----------------------|
| David Stephenson | Certified Wetland Evaluator | Project Management, |
| | Certified ELC | Reporting |
| | Certified OWES | |
| | Certified Arborist | |
| Jessica Grealey | Terrestrial and Wetland | Site Assessment |
| | Biologist | |
| | Certified ELC | |
| Tara Brenton | Terrestrial and Wetland | Site Assessment |
| | Biologist | |
| | Certified ELC | |
| | Certified OWES | |
| | Certified Arborist | |
| Charlotte Moore | Terrestrial Biologist | Site Assessment |
| Megan Pope | Terrestrial Biologist | Site Assessment, Data |
| | | Analysis, Reporting |
| Gerry Schaus | GIS Technician | Mapping |

Appendix II Vegetation Codes

Appendix II

Within Project Site and 120m boundary

OWES CLASSIFICATIONS

cS₁:

[OWES: Coniferous Swamp]

h: white birch (*Betula papyrifera*), yellow birch (*Betula alleghaniensis*), trembling aspen (*Populus tremuloides*), balsam poplar (*Populus balsamifera*)

*c: balsam fir (Abies balsamea), black spruce (Picea mariana)

dc: balsam fir (Abies balsamea)

ts: speckled alder (*Alnus incana spp. rugosa*), showy mountain-ash (*Sorbus decora*)

gc: blue-bead lily (*Clintonia borealis*), star-flower (*Trientalis borealis ssp. borealis*), bunchberry (*Cornus canadensis*), wood horsetail (*Equisetum sylvaticum*), ostrich fern (*Matteuccia struthiopteris var. pensylvanica*) m: clubmoss

cS₁₃:

[OWES: Coniferous Swamp]

*c: tamarack (*Larix laricina*), black spruce (*Picea mariana*) ts: speckled alder (*Alnus incana spp. rugosa*), showy mountain-ash (*Sorbus decora*), red-berried elderberry (*Sambucus racemosa ssp. pubens*), balsam fir (*Abies balsamea*)

ls: Labrador tea (*Ledum groenlandicum*), red raspberry (*Rubus idaeus ssp. idaeus*), red currant (*Ribes rubrum*)

gc: bracken fern (*Pteridium aquilinum var. latiusculum*), ostrich fern (*Matteuccia struthiopteris var. pensylvanica*), woodland strawberry (*Fragaria vesca ssp. americana*), bunchberry (*Cornus canadensis*), Canada mayflower (*Maianthemum canadense*) m: moss sp.

hS₈:

[OWES: Deciduous Swamp]

*h: trembling aspen (*Populus tremuloides*), white birch (*Betula papyrifera*) ls: red raspberry (*Rubus idaeus ssp. idaeus*), Canada soapberry (Shepherdia canadensis), low sweet blueberry (*Vaccinium angustifolium*), Labrador tea (*Ledum groenlandicum*)

gc: woodland strawberry (*Fragaria vesca ssp. americana*), bunchberry (*Cornus canadensis*), ostrich fern (*Matteuccia struthiopteris var. pensylvanica*), blue-bead lily (*Clintonia borealis*)

tsS_{3-5,7,18}:

[OWES: Tall Shrub Swamp]

*ts: speckled alder (Alnus incana spp. rugosa), red osier dogwood

(Cornus stolonifera)

gc: pale touch-me-not (Impatiens palidia), spinulose wood fern (*Dryopteris carthusiana*), fragrant bedstraw (*Galium triflorum*)

m: moss sp.

tsS_{11.12}:

[OWES: Tall Shrub Swamp]

*ts: speckled alder (Alnus incana spp. rugosa), Bebb's willow (Salix bebbiana)

ls: Labrador tea (Ledum groenlandicum), blueberry (Vaccinium angustifolium), Bebb's willow (Salix bebbiana), speckled alder (Alnus incana spp. rugosa)

gc: rough-leaved goldenrod (Solidago patula), Philadelphia fleabane (Erigeron philadelphicus ssp. philadelphicus), tall buttercup (Ranunculus acris)

ne: reed canary grass (Phalaris arundinacea), Bottlebrush sedge (Carex comosa), fox sedge (Carex vulpinoidea)

tsS₄₆:

[OWES: Tall Shrub Swamp]

*ts: speckled alder (Alnus incana spp. rugosa), bebb's willow (Salix bebbiana)

ls: red osier dogwood (Cornus stolonifera), red raspberry (Rubus idaeus ssp. idaeus)

gc: lady fern (Athyrium filix-femina var. angustum), tall meadowrue (Thalictrum pubescens), New England aster (Symphyotrichum novaeangliae), rough goldenrod (Solidago rugosa ssp. rugosa), Common hairgrass (Deschampia flexuosa)

ne: reed canary grass (*Phalaris arundinacea*)

neM₁₅:

[OWES: Narrow-leaved Emergents Marsh]

*ne: aquatic sedge (Carex aquatilis)

reM₁₄:

[OWES: Robust Emergents Marsh]

ds: speckled alder (Alnus incana spp. rugosa)

*re: common cattail (*Typha latifolia*)

ff: greater duckweed (Spirodela polyrhiza)

FEC CLASSIFICATIONS

ES6m: [FEC: Trembling Aspen-Black Spruce-Balsam Fir-Medium Soil] Mixedwood stands on fresh to moderately moist, medium loamy to silty soils. Medium number of shrubs, herb rich.

ELC CLASSIFICATIONS

MEGM3-8: [ELC: Reed Canary Grass Graminoid Meadow Type]

Outside of Project Site and 120m boundary

OWES CLASSIFICATIONS

cS_{2,27,32,33,34,37}:

[OWES: Coniferous Swamp]

 $tsS_{10,16,17,19-24,38,39,43-45,48,81}$:

[OWES: Tall Shrub Swamp]

 $neM_{28,40-42,83}$:

[OWES: Narrow-leaved Emergents Marsh]

reM₂₉:

[OWES: Robust Emergents Marsh]

FEC CLASSIFICATIONS

ES1r: [FEC: White Spruce-White Birch-Very Shallow Soil-Species Rich] Mixedwood dominated by white spruce and white birch on dry to fresh, very shallow soils (0-30cm) over bedrock. Medium number of shrubs, herb poor.

ES6m: [FEC: Trembling Aspen-Black Spruce-Balsam Fir-Medium Soil] Mixedwood stands on fresh to moderately moist, medium loamy to silty soils. Medium number of shrubs, herb rich. ELC CLASSIFICATIONS

THDM2-8: [ELC: Raspberry Deciduous Shrub Thicket Type] WODM5-1: [ELC: Moist Poplar Deciduous Woodland Type]

MEMM3: [ELC: Fresh Mixed Meadow Ecosite]

APPENDIX III Wetland Evaluation

| | Abitibi-Martir | n's Meadow-Empire We | tland Comp | lex | | |
|------------------------------------|-----------------------|-------------------------|------------|-----------|--------------|-----|
| | Wetland | d Evaluation Edition | | 2012 | | |
| | | February 22, 2012 | | | | |
| | | Comments | | | | |
| | | | | | | |
| | | | | | | |
| Attached Documents in | nclude: | | | | | |
| Man of Interesponding | | | | | | |
| Map of Interspersion | tland Complex Catchmo | ent Racin | | | | |
| Vascular Plant List | trana complex catemin | ent Basin | | | | |
| Fauna list | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | Additional Information | 1 | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| OCC : 1N | A.1 | 1 1 1 1 3 A 1 1 3 A 1 | F ' W | 1 10 | 1 | |
| Official Name: Evaluation Edition: | 2012 | bitibi-Martin's Meadow- | | nd ID.: | iex | |
| Wetland Significance | | Class: h Last Evaluated | wella | | 22, 2012 | |
| Provincially Significat | | h Last Updated | | reditiary | 22, 2012 | |
| Special Planning Consi | | Il Last Opdated | | | Scores | |
| Special Familia | | <u> </u> | | | Biological: | 132 |
| | | | | | Social: | 107 |
| | | | | Н | ydrological: | 205 |
| | | | | | al Features: | 159 |
| | | | | | Overall: | 603 |
| Submitted by: | | esources Solutions Inc. | - | | | |
| Date: | Fel | bruary 22, 2012 | | | | |

| ID NAME: MINISTRATIV FFICE (if differ VATION AUT nin a designated OR REGIONA HIP: CONCESSION arate sheet if ne | A VE REGION: rent from Dis HORITY JU CA, check he AL MUNICI S: eccessary) | Abitibi-Mart I: Coch strict): URISDICTIO ere: PALITY: | ON: X Cochraneyer Conc. 1 Conc. 9 Lo | -Empire Wet STRICT: Coch | Cochrane Trane nrane nc. 10 Lots 12-19, |
|--|--|---|--|---|--|
| ID NAME: MINISTRATIV FFICE (if differ VATION AUT nin a designated OR REGIONA HIP: CONCESSION arate sheet if ne | A VE REGION: rent from Dis HORITY JU CA, check he AL MUNICI S: eccessary) | Abitibi-Mart I: Coch strict): URISDICTIO ere: PALITY: | tin's Meadow nrane DIS ON: X Cochra neyer Conc. 1 Conc. 9 Lo | -Empire Wet STRICT: Coch | Cochrane |
| MINISTRATIVE FICE (if differ VATION AUT on a designated OR REGIONALIP: CONCESSION arate sheet if ne D AIR PHOTO | rent from Disconnection of the control of the contr | i: Coch strict): URISDICTIO ere: PALITY: | ON: X Cochraneyer Conc. 1 Conc. 9 Lo | Cochane 1 Lot 17, Cor | Cochrane |
| TFICE (if differ VATION AUT nin a designated OR REGIONA HIP: CONCESSION arate sheet if ne | HORITY JU: CA, check he AL MUNICI | strict): JRISDICTIO ere: PALITY: Glackm | ON: X Cochraneyer Conc. 1 Conc. 9 Lo | Cock nne 1 Lot 17, Cor | nrane |
| vation aut nin a designated OR REGIONA HIP: CONCESSION arate sheet if ne | HORITY JU CA, check he AL MUNICII | ere: | Cochra | nne 1 Lot 17, Cor | |
| OR REGIONA HIP: CONCESSION arate sheet if ne | ICA, check he AL MUNICII IS: eccessary) | ere: | Cochra | nne 1 Lot 17, Cor | |
| OR REGIONALIP: CONCESSION arate sheet if ne | AL MUNICIDES SECONDARY | Glackm | Cochra neyer Conc. 1 Conc. 9 Lo | nne 1 Lot 17, Cor | |
| CONCESSION arate sheet if ne | (S: | Glackm | neyer Conc. 1 Conc. 9 Lo | nne 1 Lot 17, Cor | |
| CONCESSION arate sheet if ne | ecessary) | | neyer Conc. 1 Conc. 9 Lo | 1 Lot 17, Cor | nc. 10 Lots 12-19, |
| arate sheet if ne | ecessary) | | Conc. 9 Lo | | nc. 10 Lots 12-19, |
| arate sheet if ne | ecessary) | | Conc. 9 Lo | | |
| | REFERENC | Conc | | | ac. 8 Lots 12-18, |
| | | CES | c. 7 Lots 13-18 | B, Conc. 6 Lo | ts 16-17, Conc. 5 Lots |
| | T 14 4 | | | | |
| | _ Longitude | e: | | | |
| reference: | | Zone: Grid:E | 17 U 501243 | | Block: N <u>5442382</u> |
| Topographic Ser | ries: | | | | |
| name(s) | | | | | |
| number(s) | | | edit | cion | |
| | | | | | |
| otographs: Date | photo taken: | Spr | ring 2005 | Scale: Go | oogle Earth Imagery |
| ate numbers | | | | | |
| ate nameers. | | | | | |
| arate sheet if ne | ecessary) | | | | |
| ase Map numbe | ers & scale | | | | |
| | | | | | |
| 1 | otographs: Date late numbers: | otographs: Date photo taken: | late numbers: arate sheet if necessary) ase Map numbers & scale | arate sheet if necessary) ase Map numbers & scale | otographs: Date photo taken: Spring 2005 Scale: Getalate numbers: arate sheet if necessary) |

| b) Wetland c | | vetland area: | - | hectares | 3 | |
|------------------------|-------------|--------------------|----------|------------|---------------|--------------|
| | omplex c | omprised of | 11 | individu | ual wetlands: | |
| Wetland U | | er | | | | Size of each |
| (for referen | nce) | | | | - | wetland unit |
| *** 1 1 ** | | W.EE 004 | Isolated | Palustrine | Riverine | Lacustrin |
| Wetland U | | WET-001 | | 33.71 | 21.00 | _ |
| Wetland U | | WET-002 | | 119.89 | 21.09 | |
| Wetland U | | WET-003 | | 9.66 | | _ |
| Wetland U | | WET-004 | | 6.09 | 01.05 | 10.04 |
| Wetland U | | WET-005 | | 277.49 | 81.35 | 10.84 |
| Wetland U | | WET-006 | | 10.97 | | |
| Wetland U | | WET-007 | | 5.19 | | |
| Wetland U | | WET-008 | | 2.03 | | |
| Wetland U | | WET-009 | | 1.53 | - | |
| Wetland U Wetland U | | WET-010 WET-011 | | 14.93 | 2.60 | |
| | | WE1-011 | | 98.15 | 3.60 | |
| Wetland U | | | | | - | |
| Wetland U Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | | | | _ |
| Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | | | | - |
| Wetland U | | | | | | |
| Wetland U | | | | | | |
| Wetland U | | | 0.00 | 579.64 | 106.04 | 10.84 |
| | | heets if necess | | 317.04 | 100.04 | 10.04 |
| (Attach au | uitionai si | neets ii necess | ary) | | | |
| | 'AI WET | LAND SIZE | | | 696.52 | ha |

1.0 BIOLOGICAL COMPONENT

1.1 PRODUCTIVITY

1.1.1 GROWING DEGREE-DAYS/SOILS

| GROV | VING DEGF | REE DAYS | S | OILS | |
|--------|-----------|-----------|---|------------|----------------|
| (check | one) | | E | stimated F | ractional Area |
| 1) | | <1600 | | 0.300 | clay/loam |
| 2) | | 1600-2000 | | | silt/marl |
| 3) | X | 2000-2400 | | | limestone |
| 4) | | 2400-2800 | | | sand |
| 5) | | 2800-3000 | | 0.200 | humic/mesic |
| 6) | | >3000 | | 0.500 | fibric |
| _ | | | | | granite |

SCORING:

| Growing | Clay- | Silt- | Lime- | Sand | Humic- | Fibric | Granite |
|-----------|-------|-------|-------|------|--------|--------|---------|
| Degree- | Loam | Marl | stone | | Mesic | | |
| Days | | | | | | | |
| <1600 | 12 | 11 | 9 | 7 | 7 | 6 | 4 |
| 1600-2000 | 15 | 13 | 11 | 9 | 8 | 7 | 5 |
| 2000-2400 | 18 | 15 | 13 | 11 | 9 | 8 | 7 |
| 2400-2800 | 22 | 18 | 15 | 13 | 11 | 9 | 7 |
| 2800-3000 | 26 | 21 | 18 | 15 | 13 | 10 | 8 |
| >3000 | 30 | 25 | 20 | 18 | 15 | 12 | 9 |

(maximum score 30; if wetland contains more than one soil type,

evaluate based on the fractional area)

Steps required for evaluation: (maximum score 30 points)

- 1. Select GDD line in evaluation table applicable to your wetland;
- 2. Determine fractional area of the wetland for each soil type;
- 3. Multiply fractional area of each soil type by score;
- 4. Sum individual soil type scores (round to nearest whole number).

In wetland complexes the evaluator should aim at determining the percentage of area occupied by the categories for the complex as a whole.

Final Score Growing Degree-Days/Soils (maximum 30 points)

| Northern Ontario Wetland Evaluation, D | Oata and Scoring Record | February 2012 |
|--|--|-----------------------------|
| 1.1.2 WETLAND TYPE (Fractional Are | a = area of wetland type/total wetland area |) |
| Fractional Area | Score | |
| Bog Fen Swamp 0.87 | x 3 x 6 x 8 0.00 0.00 6.96 | |
| Marsh 0.13 | x 15 1.95 | |
| | Wetland type score (maxi | mum 15 points) |
| 1.1.3 SITE TYPE (Fractional Area = are | ea of site type/total wetland area) | |
| | Fractional Area | Score |
| Isolated | x 1 = | 0.000 |
| Palustrine (permanent or intermittent flow) Riverine | $\begin{array}{c ccccc} $ | 1.660 |
| Riverine (at rivermouth) Lacustrine (at rivermouth Lacustrine (on enclosed | x 5 = x 5 = | 0.000 |
| bay, with barrier beach) Lacustrine (exposed to lake) | $\begin{array}{cccccccccccccccccccccccccccccccccccc$ | 0.000 0.040 |
| | Sub Total: Site Type Score (ma | 2.300 eximum 5 points) 2 |
| 1.2 BIODIVERSITY | v. | · · · · |
| 1.2.1 NUMBER OF WETLAND TYPES | | |
| (Check only one) | Score | |
| 1) one | 9 points | |
| 2) X two | 13 | |
| 3) three four | 20 30 | |
| Nu | mber of Wetland Types Score (maximum | m 30 points) 13 |
| | | |
| | | |
| | | |
| | | |
| | | |
| 1 | 4 | |

1.2.2 VEGETATION COMMUNITIES

Attach a separate sheet listing community (map) codes, vegetation forms and dominant species. Use the form on the following page to record percent area by dominant vegetation form. This information will be used in other parts of the evaluation.

Communities should be grouped by number of forms. For example, 2 form communities might appear as follows:

2 forms

| Code | Forn | ns | Don | ninant Species | _ | | |
|------|------|----|-----|------------------|-----|------------------|----------------------------|
| M6 | re, | ff | re, | Typha latifolia; | ff, | Lemna minor, | Wolffia |
| S1 | ts, | gc | ts, | Salix discolor; | gc, | lmpatiens capen. | sis, Thelypteris palustris |

Note that the dominant species for each form are separated by a semicolon. The dominant species (maximum of 2) within a form are separated by commas.

Scoring:

| Total # of communities | Total # of communities | Total # of communities |
|------------------------|-------------------------|----------------------------|
| with 1-3 forms = 40 | with $4-5$ forms = 23 | with 6 or more forms $= 1$ |
| 1 = 1.5 points | 1 = 2 points | 1 = 3 points |
| 2 = 2.5 | 2 = 3.5 | 2 = 5 |
| 3 = 3.5 | 3 = 5 | 3 = 7 |
| 4 = 4.5 | 4 = 6.5 | 4 = 9 |
| 5 = 5 | 5 = 7.5 | 5 = 10.5 |
| 6 = 5.5 | 6 = 8.5 | 6 = 12 |
| 7 = 6 | 7 = 9.5 | 7 = 13.5 |
| 8 = 6.5 | 8 = 10.5 | 8 = 15 |
| 9 = 7 | 9 = 11.5 | 9 = 16.5 |
| 10 = 7.5 | 10 = 12.5 | 10 = 18 |
| 11 = 8 | 11 = 13 | 11 = 19 |
| | | |
| +.5 each additional | +.5 each additional | + 1 each additional |
| community = 5.0 | community = 5.0 | community = 3.0 |

e.g., a wetland with 3 one form communities 8 six form communities would score:

4 two form communities

12 four form communities and

6+13.5+15=34.5=35 points

Vegetation Communities Score (maximum 45 points)

| Northern Ontario Wetland Eval | February 2012 | |
|-------------------------------|--|-----------|
| W. d. L.W. | 10 | |
| Wetland Name: | Abitibi-Martin's Meadow-Empire Wetland | 1 Complex |
| Wetland Size (ha): | 696.52 | |
| Vegetation Form | % area in which form is dominant | |
| h | 0.20 | |
| c | 30.20 | |
| dh | 0.00 | |
| de | 0.00 | |
| ts | 56.46 | |
| ls | 0.00 | |
| ds | 0.00 | |
| gc | 0.00 | |
| m | 0.00 | |
| ne | 8.82 | |
| be | 0.00 | |
| re | 4.37 | |
| ff | 0.00 | |
| f | 0.00 | |
| su | 0.00 | |
| u (unvegetated) | 0.00 | |
| Total = 100% | 100.00 | |
| | | |
| | | |
| | | |
| | | |
| | 6 | |

| Northern Ontario | Wetland Evaluation Data and Scoring Record | February 2012 |
|------------------------|---|---------------|
| 1 2 2 DIVERGITY O | E CLIDDOLINIDING HADITAT | |
| (Check all appropriate | F SURROUNDING HABITAT | |
| (Спеск ан арргориак | a ticins(1)) | |
| | recent burn (< 5 yr) | |
| | abandoned agricultural land | |
| | utility corridor | |
| X | deciduous forest | |
| | recent cutover or clearcut (<5 yr) | |
| X | coniferous forest | |
| X | mixed forest (at least 25% conifer and 75% deciduous or vice versa) | |
| X | crops | |
| | abandoned pits and quarries | |
| X | pasture | |
| | ravine | |
| X | fence rows | |
| X | open lake or deep river | |
| X | creek flood plain | |
| | rock outcrop | |
| | | |
| Div | ersity of Surrounding Habitat Score (1 for each, maximum 7 points) | 7 |
| 1 2 / DDOVIMITY T | O OTHER WETLANDS | |
| | propriate category only) | Scoring |
| (eneck mst upp | ropinuo eurogory omy) | Scoring |
| 1) x | Hydrologically connected by surface water to other wetlands | |
| | (different dominant wetland type) or open lake or river | |
| | within 1.5 km | 8 points |
| | | • |
| 2) | Hydrologically connected by surface water to other wetlands | |
| | (same dominant wetland type) within 0.5 km | 8 |
| | | |
| 3) | Hydrologically connected by surface water to other wetlands | |
| | (different dominant wetland type),or open lake or river from | |
| | 1.5 to 4 km away (Second Marsh Wetland) | 5 |
| | | |
| 4) | Hydrologically connected by surface water to other wetlands | _ |
| | (same dominant wetland type) from 0.5 to 1.5 km away | 5 |
| 5) | Wishin 0.75 land of other westlands (4:55 and 4 and and and and | |
| 5) | Within 0.75 km of other wetlands (different dominant wetland type) | |
| | or open lake or river, but not hydrologically connected by | 5 |
| | surface water | 5 |
| 6) | Within 1 km of other wetlands, but not hydrologically | |
| <u> </u> | connected by surface water | 2 |
| | connected by surface water | <u> </u> |
| 7) | No wetland within 1 km | 0 |
| Pro | eximity to other Wetlands Score (Choose one only, maximum 8 points) | 8 |
| | | |
| | 7 | |

| Northern Ontario Wetland Evaluation Data and Scoring | g Record | February 2012 |
|--|------------------------------------|---------------|
| 1.2.5 INTERSPERSION | | |
| | | |
| Number of Intersections | C | |
| (Check one) | Score | |
| 1) 26 or less | 3 | |
| 2) 27 to 40 | 6 | |
| 3) 41 to 60 | 9 | |
| 4) 61 to 80 | 12 | |
| 5) 81 to 100 | 15 | |
| 6) 101 to 125 | 18 | |
| 7) 126 to 150 | 21 | |
| 8) 151 to 175 9) 176 to 200 | 24 27 | |
| 10) >200 | 30 | |
| 10) >200 | 30 | |
| Interspersion Score (| Choose one only maximum 30 points) | 24 |
| 1.2.6 OPEN WATER TYPES | | |
| | | |
| Permanently flooded: | | |
| (Check one) | Score | |
| 1) x type 1 | 8 | |
| 2) type 1 | 8 | |
| 3) type 3 | 14 | |
| 4) type 4 | 20 | |
| 5) type 5 | 30 | |
| 6) type 6 | 8 | |
| 7) type 7 | 14 | |
| 8) type 8 | 3 | |
| 9) no open water | 0 | |
| Open Water Type Score (C | hoose one only maximum 30 points) | 8 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| 8 | | |

February 2012

1.3 SIZE

696.52 hectares 73 Subtotal for Biodiversity

Size Score (Biological Component) (maximum 50 points)

37

Evaluation Table Size Score (Biological component)

| Wetland | Total Score for Biodiversity Subcomponent | | | | | | | | | |
|-----------|---|-------|-------|-------|-------|-------|------------|-------------|-------------|------|
| size (ha) | <37 | 37-47 | 48-60 | 61-72 | 73-84 | 85-96 | 97- 108 | 109- 120 | 121- 132 | >132 |
| <20 ha | 1 | 5 | 7 | 8 | 9 | 17 | 25 | 34 | 43 | 50 |
| 20-40 | 5 | 7 | 8 | 9 | 10 | 19 | 28 | 37 | 46 | 50 |
| 41-60 | 6 | 8 | 9 | 10 | 11 | 21 | 31 | 40 | 49 | 50 |
| 61-80 | 7 | 9 | 10 | 11 | 13 | 23 | 34 | 43 | 50 | 50 |
| 81-100 | 8 | 10 | 11 | 13 | 15 | 25 | 37 | 46 | 50 | 50 |
| 101-120 | 9 | 11 | 13 | 15 | 18 | 28 | 40 | 49 | 50 | 50 |
| 121-140 | 10 | 13 | 15 | 17 | 21 | 31 | 43 | 50 | 50 | 50 |
| 141-160 | 11 | 15 | 17 | 19 | 23 | 34 | 46 | 50 | 50 | 50 |
| 161-180 | 13 | 17 | 19 | 21 | 25 | 37 | 49 | 50 | 50 | 50 |
| 181-200 | 15 | 19 | 21 | 23 | 28 | 40 | 50 | 50 | 50 | 50 |
| 201-400 | 17 | 21 | 23 | 25 | 31 | 43 | 50 | 50 | 50 | 50 |
| 401-600 | 19 | 23 | 25 | 28 | 34 | 46 | 50 | 50 | 50 | 50 |
| 601-800 | 21 | 25 | 28 | 31 | 37 | 49 | 50 | 50 | 50 | 50 |
| 801-1000 | 23 | 28 | 31 | 34 | 40 | 50 | 50 | 50 | 50 | 50 |
| 1001-1200 | 25 | 31 | 34 | 37 | 43 | 50 | 50 | 50 | 50 | 50 |
| 1201-1400 | 28 | 34 | 37 | 40 | 46 | 50 | 50 | 50 | 50 | 50 |
| 1401-1600 | 31 | 37 | 40 | 43 | 49 | 50 | 50 | 50 | 50 | 50 |
| 1601-1800 | 34 | 40 | 43 | 46 | 50 | 50 | 50 | 50 | 50 | 50 |
| 1801-2000 | 37 | 43 | 47 | 49 | 50 | 50 | 50 | 50 | 50 | 50 |
| >2000 | 40 | 46 | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 50 |

| Northern Ontario Wetland Eva | February 2012 | |
|---|--|-----------------------------|
| | 2.0 SOCIAL COMPONENT | |
| 2.1 ECONOMICALLY VALU | JABLE PRODUCTS | |
| 2.1.1 WOOD PRODUCTS | | |
| Area of wetland forested (ha), i.e. only) | dominant form is h or c. Note that this is not | t wetland size. (Check one |
| | Score | |
| 1) <5 ha | | |
| 2) 5 -25 ha | | |
| 3) 26 -50 ha | a 6 | |
| 4) 51- 100 ha | a 8 | |
| 5) 101 -200 ha | | |
| 6) X >200 ha | a 14 | |
| Source of information: | NRSI mapping | |
| | Wood Products Score (Score one only, 1 | maximum 14 points) 14 |
| | | |
| 2.1.2 Lowbush Cranberry | | Sagra (Changa ana) |
| (Check one) Present | 1) | Score (Choose one) 2 points |
| Absent | 2) 0 | 2 points 0 |
| Source of information: | | |
| | | |
| | Lowbush Cranberry Score (| (maximum 2 points) |
| 2.1.3 Wild Rice | | |
| (Check one) | | Score (Choose one) |
| Present (at least 0.5 ha) | 1) X | 10 points |
| Absent | 2) | 0 |
| Source of infolmation: | Cochrane MNR office | |
| | Wild Rice Score (maximum | 10 points) 10 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | 10 | |

| | aluation Data and | Scoring | Record | | | Februa | ry 2012 |
|---|-------------------------------------|----------|--|-------------------|----------------------------|----------|---------|
| 1.4 COMMERCIAL FISH (BA | IT FISH AND/OR | COARS | SE EISH) | | | | |
| (Check one) | II IISII AND/OK | COAK | DL 11311) | | Score (Cl | noose oi | ne) |
| Present 1) X 12 points | | | | | | | |
| Absent | 2) 0 | | | | | | |
| ource of information: | | NRSI | | | | | |
| | Con | nmercia | al Fish Score (maxin | mum 12 | points) | | 12 |
| 1.5 FURBEARERS | | | | | | | |
| (Consult Appendix 9) | | | | | | | |
| ame of furbearer | | Sourc | ee of information | | | | |
| | | | | | | | |
| beaver | 3 | | field work | | | | |
| red fox | 3 | | field work | | | | |
| red squirrel | 3 | | field work | | | | |
| marten | 3 | | Cochrane MNR | office | | | |
| | | | | | | | |
| coring: 3 points for each species | | | Furbearer Score (n | maximuı | n 12 points | s) | 12 |
| coring: 3 points for each species 2.2 RECREATIONAL ACTIV | ITIES_ | | | naximui | n 12 points | 3) | 12 |
| | ITIES_ | | Furbearer Score (n | naximui | n 12 points | s) | 12 |
| | ITIES_ | etland-A | | t/ | n 12 points Fishing | | 12 |
| 2.2 RECREATIONAL ACTIV | Type of We | etland-A | ssociated Use Nature Enjoyment | t/ | | | 12 |
| 2.2 RECREATIONAL ACTIV | Type of We Hunting 40 points 20 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 | t/ | Fishing 40 points 20 | | 12 |
| 2.2 RECREATIONAL ACTIV Intensity of Use High Moderate Low | Type of We Hunting 40 points 20 8 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 8 | t/ / | Fishing 40 points 20 8 | | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown | Type of We Hunting 40 points 20 | etland-A | SSOCIATED USE Nature Enjoyment Ecosystem Study 40 points 20 8 0 | t/ / | Fishing 40 points 20 | X | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown Totals | Type of We Hunting 40 points 20 8 0 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 8 0 | t/ / X 0 | Fishing 40 points 20 8 0 | X 8 | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown | Type of We Hunting 40 points 20 8 0 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 8 0 | t/ / X 0 | Fishing 40 points 20 8 0 | X 8 | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown Totals (score one level for each of | Type of We Hunting 40 points 20 8 0 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 8 0 | t/ / X 0 maximur | Fishing 40 points 20 8 0 | X 8 | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown Totals (score one level for each of | Type of We Hunting 40 points 20 8 0 | etland-A | ssociated Use Nature Enjoyment Ecosystem Study 40 points 20 8 0 | t/ / X 0 maximur | Fishing 40 points 20 8 0 | X 8 | 12 |
| Intensity of Use High Moderate Low Not possible/NotKnown Totals (score one level for each of | Type of We Hunting 40 points 20 8 0 | etland-A | SSOCIATED USE Nature Enjoyment Ecosystem Study 40 points 20 8 0 ores are cumulative; r | t/ X 0 maximum | Fishing 40 points 20 8 0 | X 8 | 12 |

| Northern Ontario Wetland Evaluation, Data and Scoring: R | ecord Februa | ry 2012 |
|--|--------------------------------|-----------|
| 2.3 LANDSCAPE AESTHETICS | | |
| 2.5 LANDSCAPE AESTHETICS | | |
| 2.3.1 DISTINCTNESS | | |
| (Check one) | Score (Choose one) | |
| Clearly distinct 1) | 3 points | |
| Indistinct 2) X | 0 | |
| · | | |
| Landscape Distinctn | ess Score (maximum 3 points) | 0 |
| 2.3.2 ABSENCE OF HUMAN DISTURBANCE | | |
| (Charle and) | Sagra (Changa ana) | |
| (Check one) Human disturbances absent or nearly so | Score (Choose one) 7 points | |
| One or several localized disturbances | 2) X 4 | |
| Moderate disturbance; localized water pollution | 3) 2 | |
| Wetland intact but impairment of ecosystem quality | 3)2 | |
| intense in some areas | 4) 1 | |
| Extreme ecological degradation, or water pollution | , | |
| severe and widespread | 5) 0 | |
| Source of information: air photo | s, field work | |
| | | |
| Absence of Human Distur | bance Score (maximum 7 points) | 4 |
| A A EDVICATION AND DUDI IC AWA DENIEGO | | |
| 2.4 EDUCATION AND PUBLIC AWARENESS | | |
| 2.4.1 EDUCATIONAL USES | | |
| (Check one) | Score (Choose one) | |
| Frequent 1) | 20 points | |
| Infrequent 2) | 12 | |
| No visits 3) X | 0 | |
| 3) | O . | |
| Source of information: Cochr | ane MNR office | |
| | | |
| Educational Us | ses Score (maximum 20 points) | 0 |
| 2.4.2 FACILITIES AND PROGRAMS | | |
| | | |
| (check one) | Score (Che | oose one) |
| Staffed interpretation centre | 1) 8 points | |
| No interpretation centre or staff but a system of | <u> </u> | |
| self-guiding trails or brochures available | 2) 4 | |
| Facilities such as maintained paths (e.g., woodchips) | · <u></u> | |
| boardwalks, boat launches or observation towers | | |
| but no brochures or other interpretation | 3)2 | |
| No facilities or programs | 4) X 0 | |
| Source of information: Cochr | ane MNR office | |
| | | |
| Facilities and Progra | ms Score (maximum 8 points) | 0 |

| Northern Ontario Wetland Evaluation, Data and Scoring Record February 2012 | | | | | | | |
|---|--|---------|--------|------------------------|--------------|------------------------------|----|
| 2.4.3 RESEARCH AND STUDIES | | | | | | | |
| 2.4.3 RESEARCH AND STUDIES (check appropriate spaces) Long term research has been done Research papers published in refere journal or as a thesis One or more (non-research) reports on some aspect of the wetland 's flo hydrology etc. No research or reports Attach list of known reports by above | | X | | Score 12 points 10 5 0 | | | |
| Research and St | udies Score (Scor | e is cu | mula | tive, maxim | um 12 | 2 points) | 0 |
| 2.5 PROXIMITY TO AREAS OF H | IIMAN SETTI EN | MENT | ין | | | | |
| Circle the highest applicable score | UNIAN SETTLE | VIIVI | L | - | | | |
| Distance of wetland from | 1) | | 2) | populati | | 3) population | |
| settlement | population> 10 | ,000 | | 2,500 -10 | ,000 | <2,500 or cotta community | _ |
| 1) Within or adjoining | 40 points | | | 26 | | 16 | |
| settlement | | | | | | | |
| 2) 0.5 to 10 km from settlement | 26 | | | 16 | X | 10 | |
| 3) 10 to 60 km from settlement | 12 | | | 8 | | 4 | |
| 4) >60 km from settlement | 5 | | | 2 | | 0 | |
| 5) >100 km from settlement | 0 | 0 | | 0 | 16 | 0 | 0 |
| | | U | | | 10 | | U |
| Name of settlement: | Town | of Co | chrane | e | | | |
| Proxi | imity to Human S | ettlen | nent S | core (maxin | num 4 | 10 points) | .6 |
| 2.6 OWNERSHIP (FA= fraction Are | - | | | · · | | Score | |
| 2.0 OWNERSHII (FA- Haction Arc |) | | | | | Score | |
| FA of wetland in public or private on held under contract or in trust for we FA of wetland area in public owners FA of wetland area in private owners. | etland protection ship,not as above | | 1. | x x 00 x | 10 8 4 | = 0.00 = 0.00 = 4.00 | |
| Source of information: | Cochrar | ne MN | R offi | ice | | | |
| | | Own | ershin | Score (max | kimun | 1 10 points) | 4 |
| | | | г | | | | |
| | | | | | | | |
| | | | | | | | |
| | 13 | | | | | | |

February 2012

2.7 SIZE

696.52 hectares 80 Subtotal for Social

Evaluation Table for Size Score (Social Component)

| | rable | ioi size sco | re (Social C | omponent) | | | | | | |
|----------------------|--------------------------------|--------------|--------------|-----------|-------|--------|---------|---------|---------|------|
| Wetland Size (ha) | Lotal for Size Dependent Score | | | | | | | | | |
| Size (iiii) | <31 | 31-45 | 46-60 | 61-75 | 76-90 | 91-105 | 106-109 | 121-135 | 136-150 | >150 |
| <2 ha | 1 | 2 | 4 | 8 | 10 | 12 | 14 | 14 | 14 | 15 |
| 2 - 4ha | 1 | 2 | 4 | 8 | 12 | 13 | 14 | 14 | 15 | 16 |
| 5 - 8ha | 2 | 2 | 5 | 9 | 13 | 14 | 15 | 15 | 16 | 16 |
| 9 - 12ha | 3 | 3 | 6 | 10 | 14 | 15 | 15 | 16 | 17 | 17 |
| 13-17 | 3 | 4 | 7 | 10 | 14 | 15 | 16 | 16 | 17 | 17 |
| 18-28 | 4 | 5 | 8 | 11 | 15 | 16 | 16 | 17 | 17 | 18 |
| 29-37 | 5 | 7 | 10 | 13 | 16 | 17 | 18 | 18 | 19 | 19 |
| 38-49 | 5 | 7 | 10 | 13 | 16 | 17 | 18 | 18 | 19 | 20 |
| 50-62 | 5 | 8 | 11 | 14 | 17 | 17 | 18 | 19 | 20 | 20 |
| 63-81 | 5 | 8 | 11 | 15 | 17 | 18 | 19 | 20 | 20 | 20 |
| 82-105 | 6 | 9 | 11 | 15 | 18 | 18 | 19 | 20 | 20 | 20 |
| 106-137 | 6 | 9 | 12 | 16 | 18 | 19 | 20 | 20 | 20 | 20 |
| 138-178 | 6 | 9 | 13 | 16 | 18 | 19 | 20 | 20 | 20 | 20 |
| 179-233 | 6 | 9 | 13 | 16 | 18 | 20 | 20 | 20 | 20 | 20 |
| 234-302 | 7 | 9 | 13 | 16 | 18 | 20 | 20 | 20 | 20 | 20 |
| 303-393 | 7 | 9 | 14 | 17 | 18 | 20 | 20 | 20 | 20 | 20 |
| 394-511 | 7 | 10 | 14 | 17 | 18 | 20 | 20 | 20 | 20 | 20 |
| 512-665 | 7 | 10 | 14 | 17 | 18 | 20 | 20 | 20 | 20 | 20 |
| 666-863 | 7 | 10 | 14 | 17 | 19 | 20 | 20 | 20 | 20 | 20 |
| 864-1123 | 8 | 12 | 15 | 17 | 19 | 20 | 20 | 20 | 20 | 20 |
| 1124-1460 | 8 | 12 | 15 | 17 | 19 | 20 | 20 | 20 | 20 | 20 |
| 1461-1898 | 8 | 13 | 15 | 18 | 19 | 20 | 20 | 20 | 20 | 20 |
| 1899-2467 | 8 | 14 | 16 | 18 | 20 | 20 | 20 | 20 | 20 | 20 |
| >2467 | 8 | 14 | 16 | 18 | 20 | 20 | 20 | 20 | 20 | 20 |

Total Size Score (Social Component)

Northern Ontario Wetland Evaluation, Data and Scoring Record February 2012 ABORIGINAL AND CULTURAL HERITAGE VALUES Either or both Aboriginal or Cultural Values may be scored. However, the maximum score permitted for 2.8 is 30 points. Attach documentation. 2.8.1 ABORIGINAL VALUES Full documentation of sources must be attached to the data record. 1) Significant 30 points 2) Not Significant 0 3) Unknown 0 Total: 2.8.2 CULTURAL HERITAGE 30 points 1) Significant 2) Not Significant 0 3) 0 Unknown Total: Aboriginal Values/Cultural Heritage Score (maximum 30 points)

February 2012

3.0 HYDROLOGICAL COMPONENT

3.1 FLOOD ATTENUATION

If the wetland is a complex including isolated wetlands, apportion the l00 points according to area. For example if 10 ha of a l00 ha complex is isolated, the isolated portion receives the maximum proportional score of 10. The remainder of the wetland is then evaluated out of 90.

Step 1: If wetland is entirely <u>Isolated</u>, go directly to Step 5.

If wetland is lacustrine and the ratio of wetland area: lake area is <0.1, or wetland is

riverine on the St. Mary's River, go to Step 5

All other wetlands, go through steps 2, 3, 4 and 5.

Step 2: Determination of Upstream Detention Factor (DF)

| (a) | Wetland area (ha) | | | 696.52 |
|-----|--|--------|--|--------|
| (b) | Total area (ha) of upstream detention area | 710.96 | | |
| | (include the wetland itself) | | | |
| (c) | Ratio of (a):(b) | | | 0.98 |
| (d) | Upstream detention factor: (c) x 2 = | 1.96 | | 1.00 |
| | (maximum allowable factor = 1) | | | |

Step 3: Determination of Peak Flow Attenuation Factor (AF)

| (a) | Wetland area (ha) | | | 696.52 |
|-----|--|---------|--|--------|
| (b) | Size of catchment basin (ha) upstream of | wetland | | |
| | (include wetland itself in catchment area) | 2198.44 | | |
| (c) | Ratio of (a):(b) | | | 0.32 |
| (d) | Wetland attenuation factor: (c) x 10 = | 3.2 | | 1.00 |
| | (maximum allowable factor = 1) | | | |

Step 4: Determination of Wetland Surface Form Factor (FF)

From the list below, select the surface form which best describes the wetland.

| | Factor | |
|--|--------|-----|
| Flooded with little or no aquatic vegetation | | 0 |
| Flooded but with submergent, emergent or floating vegetation | | 0.2 |
| Flat (lawn) vegetation (typical of fens) | | 0.5 |
| Hummock-depression microtopography | X | 0.7 |
| Patterned (e.g., string bog, ribbed fen) | | 1 |
| Surface Form Factor (FF) | 0.7 | |

(Maximum allowable factor = 1)

February 2012

Step 5:

1. Wetland is entirely Isolated 100 points

2. Wetland is lacustrine and the ratio of 0 points

wetland area: lake area is <0.1

3. Wetland is riverine along the St. Mary's River 0 points

4. For all other wetlands*, calculate as follows:

Upstream Detention Factor (DF) (Step 2) 1.00 Wetland Attenuation Factor (AF) (Step 3) 1.00 c) Surface Form Factor (FF) (Step 4) 0.70

> $[(DF + AF + FF)/3] \times 100*$ 90

*Unless wetland is a complex including isolated portions -- see above

Total Flood Attenuation Score (maximum 100 points)

0.00

GROUND WATER RECHARGE

3.2.1 SITE TYPE

Wetland > 50% lacustrine (by area) or located on the (a)

St. Mary's River

FA of lacustrine wetland (wetland <50% lacustrine)

Score = 0Wetland not as above. Calculate final score as follows:

(b) (FA= area of site type/total area of wetland)

0.83 FA of isolated or palustrine wetland 16.60 20 0.15 FA of riverine wetland 5 0.75

Site Type Score: (maximum 20 points)

3.2.2 SOILS

EVALUATION:

0.02

| Dominant Wetland Type | Sand, loam, gravel, till | | Clay or bedrock | |
|------------------------------------|--------------------------|---|-----------------|---|
| Lacustrine or on St. Mary's River | 0 | | 0 | |
| Isolated | 10 | | 5 | |
| Palustrine | 7 | X | 4 | |
| Riverine (not on St. Mary's River) | 5 | | 2 | |
| Totals | | 7 | | 0 |

Hydrological Soil Class Score (maximum 10 points)

February 2012

3.3 DOWNSTREAM WATER QUALITY IMPROVEMENT

3.3.1 WATERSHED IMPROVEMENT FACTOR

Calculation of Watershed Improvement Score is based upon the fractional area (FA) of each site type within the wetland. FA = area of site type/total area of the wetland.

| Site Type | <u>Impro</u> | vement Fac | tor (IF | <u>)</u> | |
|--------------------------------------|--------------|------------|---------|----------|-------|
| Isolated | FA | 0 | X | 0.5 = | 0.00 |
| Riverine | FA | 0.15 | X | 1 = | 0.15 |
| Palustrine with no inflow | FA | 0 | X | 0.7 = | 0.00 |
| Palustrine with inflows | FA | 0.83 | X | 1 = | 0.83 |
| Lacustrine on lake shoreline | FA | 0.02 | X | 0.2 = | 0.004 |
| Lacustrine at lake inflow or outflow | FA | 0 | X | 1 = | 0.00 |

Watershed Improvement Score (IF x 30) (maximum = 30)

20.52

3.3.2 ADJACENT AND WATERSHED LAND USE

EVALUATION

Step 1: Determination of Maximum Initial Score

Wetland on the Great Lakes or St. Mary's River (Go to Step 5a)

X All other wetlands (Go through steps 2, 3,4 and 5b)

Step 2: Determination of Broad Upslope Land Use (BLU)

Assess broad upslope land uses within the previous 5 years, agriculture, or other activities which alter the natural vegetation cover in an extensive manner.

| Choose one | | Score |
|---------------------------|---|-------|
| >50% of catchment basin | | 20 |
| 20-50% of catchment basin | | 14 |
| <20% of catchment basin | X | 4 |

Score for BLU

4

Step 3: Determination of Linear Upslope Land Uses (LUU)

Assess linear upslope uses (LUU) e.g., roads, railways, hydro corridors, pipelines, etc., crossing the upslope catchment within 200m of the wetland boundary.

| Choose the highest only | | Score |
|-------------------------|---|-------|
| Major corridor* | | 15 |
| Secondary corridor | | 11 |
| Tertiary corridor | X | 6 |
| Temporary or abandoned | | 3 |
| None | | 0 |

Score for LUU

6

Major, secondary and tertiary roads are those that are indicated as such on the provincial highways maps. Major hydro corridors are trunk lines coming directly from a generating station. Major pipelines are transcontinental lines. Secondary corridors are regional distribution lines (i.e. multi-cable hydro corridors not emanating directly from a generating station or regional gas distribution lines). Tertiary corridors are single hydro lines or local gas distribution lines (i.e. to domestic users).

| | Present 15 Not present X 0 | Score for PS | 0 | | |
|---------------|--|------------------------|-------------------|-------------------|---|
| <u>tep 5:</u> | Calculation of total score for A | djacent and Watershed | Land Use | | |
| | Wetland on the Great Lakes or St. M. All other wetlands, calculate as follows: | | | | |
| | | Final Score Bl | LU+LUU+P\$ | 10 | |
| .3.3 VE | EGETATION FORM | | | | |
| | oose the category that best describes getation of the wetland | s the | | | |
| Em | tees, shrubs or herbs (h, c, ts, ls, gc) hergents, submergents (ne, re, be, f, tele or no vegetation (u) | ff, su) | Sco 8 pc 10 | re pints | |
| 3.4 | Do CARBON SINK | ominant Vegetation For | m Score (ma | eximum 10 points) | 8 |
| Che | oose the category that best describes | s the wetland | | | |
| 1) | Wetland a bog or fen with >50% | organic soils | | 15 points | |
| 2) | Wetland has organic soils occupy of the area (i.e. mainly mineral o | = | | 6 | |
| 2) | soils, any wetland type) | | | | |
| 3) | soils, any wetland type) Marshes and swamps with >50% | organic soil | X | 9 | |
| | | - | X | 9 | |
| 3) | Marshes and swamps with >50% | oils organic | | | 9 |
| 3) | Marshes and swamps with >50% | oils organic | | 0 | 9 |
| 3) | Marshes and swamps with >50% | oils organic | | 0 | 9 |
| 3) | Marshes and swamps with >50% | oils organic | | 0 | 9 |

Northern Ontario Wetland Evaluation

February 2012

| Northern | Ontario | Wetland | Eval | nation |
|----------|---------|----------|------|--------|
| Normen | Omano | vv Cuanu | Lvai | uauon |

February 2012

3.5 SHORELINE EROSION CONTROL

From the wetland vegetation map determine the <u>dominant</u> vegetation type within the erosion zone for <u>lacustrine and riverine site type areas only.</u> Score according to the factors listed below.

Step 1:

Score

0

Wetland entirely isolated or palustrine

Any part of the Wetland riverine or lacustrine (proceed to Step 2)

Step 2:

Choose the one characteristic that best describes the shoreline vegetation (see text for a definition of shoreline)

| | | | Score |
|----|---|----------------------------|-------|
| 1) | | Trees and shrubs | 15 |
| 2) | X | Emergent vegetation | 8 |
| 3) | | Submergent vegetation | 6 |
| 4) | | Other shoreline vegetation | 3 |
| 5) | | No vegetation | 0 |

Shoreline Erosion Control Score (maximum 15 points)

8

3.6 GROUNDWATER DISCHARGE

(Circle the characteristics that best describe the wetland being evaluated and then sum the scores)

| Category | | C | Catchment Interaction | | | |
|-----------------------|--------------------|---|------------------------|---|----------------------|----|
| Wetland type | Bog = 0 | | Swamp/Marsh = 2 | 2 | Fen = 5 | |
| Basin topography | Flat/Rolling = 5 | | Hilly = 2 | | Major relief | |
| | | 5 | | | break = 5 | |
| Wetland area: Upslope | Large (>50%) = 0 | | Moderate | | Small ($<5\%$) = 5 | |
| catchment area | | | (6-50%) = 2 | 2 | | |
| Lagg Development | None found $= 0$ | 0 | Minor = 2 | | Extensive $= 5$ | |
| Seeps at wetland | None found $= 0$ | | 1-3 seeps = 5 | | 4 or more | |
| edge | | 0 | | | seeps = 10 | |
| Iron precipitates | None = 0 | | 1-3 deposits = 2 | | 4 or more | |
| evident at edge | | 0 | | | deposits = 5 | |
| Surface marl deposits | None = 0 | 0 | 1-3 deposits = 2 | | >3 = 5 | |
| Wetland pH | Low < 4.2 = 0 | | Moderate $4.2-5.7 = 5$ | | High $> 5.7 = 10$ | 10 |
| Catchment soil | Patchy = 0 | | Thin $(<20cm) = 2$ | | Thick $= 5$ | |
| coverage | | | | | | 5 |
| Catchment soil | Low = 0 | | Moderate = 2 | | High = 5 | |
| permeability | | | | 2 | | |
| Totals | | 5 | | 6 | | 15 |

(Scores are cumulative maximum score 30 points)

Groundwater Discharge Score (maximum 30 points)

| Northern Ontario Wetland Evaluation Data and Scoring Reco |
|---|
|---|

February 2012

4.0 SPECIAL FEATURES COMPONENT

4.1 RARITY

4.1.1 WETLANDS

Hills Site Region and Site District (5E only):
Wetland type (check one or more)

| | Bog |
|---|-------|
| | Fen |
| X | Swamp |
| X | Marsh |

Evaluation Table for Scoring Rarity of Wetland Type.

| Unit | Site Region | | _ | _ | _ |
|--------|-------------------|-------|-------|-----|-----|
| Number | & District | Marsh | Swamp | Fen | Bog |
| 2E | James Bay | 20 | 20 | 0 | 20 |
| 2W | Big Trout Lake | 20 | 20 | 0 | 10 |
| 3E | Lake Abitibi | 20 | 20 | 10 | 0 |
| 3W | Lake Nipigon | 20 | 20 | 10 | 0 |
| 3S | Lake St. Joseph | 20 | 20 | 10 | 0 |
| 4E | Lake Temagami | 20 | 20 | 10 | 0 |
| 4W | Pigeon River | 20 | 10 | 20 | 0 |
| 4S | Wabigoon Lake | 20 | 10 | 20 | 0 |
| 5E-1 | Thessalon | 10 | 0 | 30 | 20 |
| 5E-2 | Gore Bay | 20 | 0 | 20 | 20 |
| 5E-3 | La Cloche | 20 | 0 | 30 | 20 |
| 5E-4 | Sudbury | 10 | 0 | 30 | 10 |
| 5E-5 | North Bay | 10 | 0 | 20 | 0 |
| 5E-6 | Tomiko | 10 | 0 | 20 | 0 |
| 5E-7 | Parry Sound | 20 | 0 | 30 | 20 |
| 5E-8 | Huntsville | 20 | 0 | 30 | 20 |
| 5E-9 | Algonquin Park | 10 | 0 | 30 | 0 |
| 5E-10 | Brent | 20 | 0 | 30 | 0 |
| 5E-11 | Bancroft | 0 | 10 | 30 | 10 |
| 5E-12 | Renfrew | 0 | 0 | 30 | 10 |
| 5E-13 | Batchewana | 10 | 0 | 10 | 30 |
| 5-S | Lake of the Woods | 10 | 10 | 20 | 10 |

Rarity of Wetland Type Score (maximum 70 points)

40

| Northern Ontario Wetland Evaluation | , Data and Scoring Record | Fel | oruary 2012 |
|--|----------------------------|-----------------------|-------------|
| 3.1.2 SPECIES | | | |
| 4.1.2.1 BREEDING HABITAT | FOR AN ENDANGERE | D SPECIES | |
| Name of species | | Source of information | |
| - | | ¬ | |
| 1) | | | |
| 2) | | | |
| 4) | | | |
| 5) | | | |
| Total: | 0 | | |
| ttach documentation. | | _ | |
| ooring: | | | |
| coring: For one species | 250 points | | |
| For each additional species | 250 points | | |
| core is cumulative, no maximum score) | | | |
| Breeding Habitat for En | ndangered Species Score (1 | no maximum) | 0 |
| 4.1.2.2 TRADITIONAL MIGRATIONAL | ON OR FEEDING HABIT | TAT FOR AN ENDANGERE | D SPECIES |
| | | | |
| Name of species 1) | | Source of information | |
| 2) | | | |
| 3) | | | |
| 4) | | | |
| 5) | | | |
| Total: | 0 | | |
| ttach documentation. | | | |
| coring: | | | |
| For one species For each additional species | 150 points 75 | | |
| core is cumulative, no maximum score) | | | |
| Traditional Habita | at for Endangered Species | Score (no maximum) | 0 |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

| 1) 2) 3) 4) 5) 6) 7) 8) 9) 11) 12) | Name of | species | | | J. Sch | Source of information | n |
|--|------------|------------|------------------|--------------------------|--------------|-------------------------|----------|
| 2) 3) 4) 5) 6) 7) 8) 9) 10) 11) | | | | | | Source of information | n |
| 2) 3) 4) 5) 6) 7) 8) 9) 10) 11) | | | | | | | |
| 3) 4) 5) 6) 7) 8) 9) 10) 11) | | | | | | | |
| 4) 5) 6) 7) 8) 9) 10) 11) | | | | | | | |
| 5) 6) 7) 8) 9) 10) 11) | | | | | | | |
| 6) 7) 8) 9) 10) 11) | | | | | | | |
| 7) 8) 9) 10) 11) | | | | | | | |
| 9) 10) 11) | | | | | | | |
| 10) 11) | | | | | | | |
| 11) | | | | | | | |
| | | | | _ | | | |
| 141 | • | | | | | | |
| 13) | | | | | | | |
| 14) | | | | | | - | |
| 15) | | | | | | | |
| , | Attach s | eparate li | ist if necessary | ; Attach documenta | ition | | |
| | | | 50 points | pecies in the wetlar | | 154 | |
| 1 spe 2 spe | | = | 80 | 14 species 15 species | = | 156 | |
| 3 spe | | = | 95 | 16 species | = | 158 | |
| 4 spe | | = | 105 | 17 species | = | 160 | |
| 5 spe | cies | = | 115 | 18 species | = | 162 | |
| 6 spe | | = | 125 | 19 species | = | 164 | |
| 7 spe | | = | 130 | 20 species | = | 166 | |
| 8 spe | | = | 135 | 21 species | = | 168 | |
| 9 spe | | = | 140 | 22 species | = | 170 | |
| 0 spe | | = | 143 | 23 species | = | 172 | |
| 1 spe | | = | 146 | 24 species | _ = | 174 176 | |
| 12 spe 13 spe | | = | 149 152 | 25 species | = | 170 | |
| _ | | | | example, 26 specie | $es = 177_1$ | points, 27 species = 17 | 78 |
| nts etc. |) | | | - | | • | |
| maxin | num score) | | | | | | |
| | | | Provin | cially Significant A | Animal S | pecies Score (no max | cimum) (|
| | | | | | | _ | |
| | | | | | | | |

| (Scientific Common 1) | e names must be Name | Sc | mentation | n | Source | e of information |
|--|-------------------------|----------------------|-----------|-------------------|---------------|------------------|
| Common 1) 2) 3) 4) 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separates a conting: mber of provincially a conting: meetics = conting: cont | Name | essary; Attach docu | mentation | n | Source | e of information |
| 2) 3) 4) 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach seporing: mber of provincially pecies = peci | parate list if nec | essary; Attach docu | etland: | | | |
| 2) 3) 4) 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separations: mber of provincially species = speci | parate list if nec | essary; Attach docu | etland: | | | |
| 3) 4) 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separates services = ser | parate list if nec | essary; Attach docu | etland: | | | |
| 4) 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separates = secies = seci | parate list if nec | essary; Attach docu | etland: | | | |
| 5) 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separing: mber of provincially ecies = | parate list if nec | essary; Attach docu | etland: | | | |
| 6) 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separing: mber of provincially secies = | parate list if nec | essary; Attach docu | etland: | | | |
| 7) 8) 9) 10) 11) 12) 13) 14) 15) Attach separing: mber of provincially species = spe | parate list if nec | essary; Attach docu | etland: | | | |
| 8) 9) 10) 11) 12) 13) 14) 15) Attach separing: mber of provincially species = specie | parate list if nec | essary; Attach docu | etland: | | | |
| 9) 10) 11) 12) 13) 14) 15) Attach separates are cies are | parate list if nec | essary; Attach docu | etland: | | | |
| 10) 11) 12) 13) 14) 15) Attach separates are cies = cies | parate list if nec | essary; Attach docu | etland: | | | |
| 11) 12) 13) 14) 15) Attach separing: mber of provincially species = specie | parate list if nec | essary; Attach docu | etland: | | | |
| 12) 13) 14) 15) Attach separing: mber of provincially species = | parate list if nec | essary; Attach docu | etland: | | | |
| Attach separing: The provincial species = spe | parate list if nec | essary; Attach docu | etland: | | | |
| Attach separations: Attach separations: The provincial separation of provincial separation of separations: The provincial separation of separation of separations: The provincial separation of se | parate list if nec | essary; Attach docu | etland: | | | |
| Attach seporting: mber of provincially species = specie | parate list if nec | essary; Attach docu | etland: | | | |
| ring: mber of provinciall pecies = | | · | etland: | | | |
| pecies = | | | | | | |
| pecies = | 50 points | 14 species | = | 154 | | |
| pecies = pec | 80 | 15 species | = | 156 | | |
| pecies = pec | 95 | 16 species | = | 158 | | |
| pecies = pec | 105 | 17 species | = | 160 | | |
| pecies = pecies = pecies = | 115 | 18 species | = | 162 | | |
| pecies = pecies = | 125 | 19 species | = | 164 | | |
| pecies = | 130 | 20 species | = | 166 | | |
| | 135 | 21 species | = | 168 | | |
| species – | 140 | 22 species | = | 170 | | |
| - | 143 | 23 species | = | 172 | | |
| species = | 146 | 24 species | = | 174 | | |
| species = | 149 | 25 species | = | 176 | | |
| species = | 152 | | | | | |
| d one point for events etc.) | ry species past 2 | 25 (for example, 26 | species = | = 177 points, 27 | species = 178 | |
| | | | ~ | | | |
| | - . | cially Significant P | 'lant Spe | ecies Score (no r | maximum) | |

| Northern Ontario Wetland Evaluation, Data and Scoring Reco |
|--|
|--|

February 2012

4.1.2.5 REGIONALLY SIGNIFICANT SPECIES (SITE REGION)

Scientific names must be recorded for plant species. Lists of significant species must be approved by MNR.

SIGNIFICANT IN SITE REGION:

| C | Common Name | Scientific Name | Source of information | |
|-----|-------------------|------------------------|-----------------------|--|
| 1) | eastern phoebe | Sayornis phoebe | NRSI field work | |
| 2) | gray catbird | Dumetella carolinensis | NRSI field work | |
| 3) | northern cardinal | Cardinalis cardinalis | NRSI field work | |
| 4) | sandhill crane | Grus canadensis | NRSI field work | |
| 5) | scarlet tanager | Piranga olivacea | NRSI field work | |
| 6) | | | | |
| 7) | | | | |
| 8) | | | | |
| 9) | | | | |
| 10) | | | · | |
| 11) | | | · | |
| 12) | | | · • | |
| 13) | | | | |
| 14) | | | | |
| 15) | | | | |

Attach separate list if necessary. Attach documentation.

No. of species significant in Site Region

| 1 species | = | 20 | 6 species | = | 55 |
|-----------|---|----|------------|---|----|
| 2 species | = | 30 | 7 species | = | 58 |
| 3 species | = | 40 | 8 species | = | 61 |
| 4 species | = | 45 | 9 species | = | 64 |
| 5 species | = | 50 | 10 species | = | 67 |

Add one point for every species past 10 (no maximum score).

Significant Species (Site Region) Score (no maximum)

^{**} Score only if there is an approved list Scoring:

| Northern | Ontario | Wetland | Evaluation, | Data an | d Scoring | Record |
|----------|---------|---------|-------------|---------|-----------|--------|
| | | | | | | |

February 2012

4.2.1.6 LOCALLY SIGNIFICANT SPECIES (SITE DISTRICT)

Scientific names must be recorded for plant species. Lists of significant species must be approved by MNR.

| Common Name | Scientific Name | Source of information |
|-------------|-----------------|-----------------------|
| | | |
| 1 | | |
| 2 | | |
| 3 | | |
| 4 | | |
| 5 | | |
| 6 | | |
| 7 | | |
| 8 | | |
| 9 | | |
| 10 | | |
| 11 | | |
| 12 | | |
| 13 | | |
| 14 | | |
| 15 | <u> </u> | |
| 16 | <u> </u> | |
| 17 | <u> </u> | |
| 18 | | |

Attach separate list if necessary. Attach documentation.

Scoring:

No. of species significant in Site District

| 1 species | = | 10 | 6 species | = | 41 |
|-----------|---|----|------------|---|----|
| 2 species | = | 17 | 7 species | = | 43 |
| 3 species | = | 24 | 8 species | = | 45 |
| 4 species | = | 31 | 9 species | = | 47 |
| 5 species | = | 38 | 10 species | = | 49 |

For each significant species over 10 in the wetland, add 1 point.

Locally Significant Species (Site District) Score (no maximum)

Λ

Northern Ontario Wetland Evaluation February 2012 4.1.2.7 SPECIES OF SPECIAL STATUS Black Duck Suitable breeding habitat present and within assessment range (Figure 17) Assessment Category Check one Score 40-80 Indicated Pairs/100 km sq 25 points 20-40 Indicated Pairs/100 km sq 20 10-20 Indicated Pairs/100 km sq X 15 5-10 Indicated Pairs/100 km sq 10 1-5 Indicated Pairs/100 km sq 5 0 Habitat not suitable 0 Out of assessment range Black Duck Score (maximum 25 points) 4.2 SIGNIFICANT FEATURES AND/OR FISH & WILDLIFE HABITAT 4.2.1 NESTING OF COLONIAL WATERBIRDS Status Name of species Source of Information Score Currently nesting 50 points Known to have nested 25 within past 5 years Active feeding area (great blue heron excluded) 15 None known X 0 Attach documentation (nest locations etc., if known) Colonial Waterbirds Score (maximum 50 points) 4.2.2. WINTER COVER FOR WILDLIFE (Check only highest level of significance) Score (one only) Provincially significant 100 1) 2) Significant in Site Region 50 Significant in Site District 25 3) 3) Locally significant 10 Little or poor winter cover present 0 4) Source of information: Winter Cover for Wildlife Score (maximum 100 points) 0

| Northe | rn Ontario Wetland Evaluation | , Data and S | Scoring Record | | Febr | uary 2012 |
|-------------|--|--------------|------------------|-------------------|---------------|-----------|
| | | | | | | |
| 4.2.3 WA | TERFOWL STAGING AND/O | OR MOULT | ING | | | |
| (Check on | ly highest level of significance | for both sta | ging and moultin | ar score is cum | ulative | |
| | umns, maximum score 150) | 101 bour su | gilig and moutan | g, score is cuiii | uiative | |
| | , in | | | | | |
| | | Staging | Score | Moulting | Score | |
| | | | (one only) | | (one only) | |
| 1) | Nationally significant | | 150 | | 150 | |
| 2) | Provincially significant | | 100 | | 100 | |
| 3) | Regionally significant | | 50 | | 50 | |
| 4) | Known to occur | | 10 | | 10 | |
| 5) | Not possible | T/ | 0 | N/ | 0 | |
| 6) | Not known | X | 0 | X | 0 | |
| | Total: | | _ | 0 | | |
| Source of i | information: | | | | | |
| Bource of | | l Moulting | and Staging Sco | re (maximum | 150 points) | 0 |
| | | | | | , | |
| 4.2.4 WA | TERFOWL BREEDING | | | | | |
| | | _ | | | | |
| | (Check only highest level of | significance |) Sco | ore | | |
| | | | | | | |
| 1) | Provincially sign | | | 00 | | |
| 2) | Regionally signif | ficant | | 50 | | |
| 3) | X Habitat suitable | 11. | 1 | 10 | | |
| 4) | Habitat not suital | ble | | 0 | | |
| Source of i | information: | | field work | | | |
| | | | | | | |
| | | Waterfow | l Breeding Score | e (maximum l0 | OO points) | 10 |
| | | 05 | amo | | | |
| 4.2.5 MIC | GRATOR PASSERINE, SHOI | REBIRD OF | RAPTOR STO | POVER AREA | | |
| | (abaak highast annliaghle cat | 22041) | | | | |
| | (check highest applicable cat | egory) | | | | |
| 1) | Provincially sign | ificant | 10 | 00 | | |
| 2) | Significant in Sit | | | 50 | | |
| 3) | Significant in Sit | | | 10 | | |
| 4) | X Not significant | | | 0 | | |
| | | | | | | |
| Source of | information: | | | | | |
| | | | | | | |
| | Passerine, Shore | ebird or Ra | ptor Stopover S | core (maximur | n 100 points) | 0 |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | 28 | | | |

| Northern Ontario Wetland Evaluation, Data and Scoring Reco | ord | February 2012 |
|---|-------------------------------|---------------|
| 4.2.6 UNGULATE HABITAT | | |
| EVALUATION | | |
| | | |
| Score $(1) + (2) + $ one of (3) to (6) | C | |
| (1) V Un gulete summer sever | Score | |
| (1) X Ungulate summer cover (2) Mineral licks | 15 points 50 | |
| (2) willer at ficks | 30 | |
| (3) Moose aquatic feeding area Class 1 | 0 | |
| (4) X Moose aquatic feeding area Class 2 | 10 | |
| (5) Moose aquatic feeding area Class 3 | 20 | |
| (6) Moose aquatic feeding area Class 4 | 35 | |
| Score is cumulative for a maximum possible score of 100) | | |
| | re (maximum 100 points) | 25 |
| | | |
| | | |
| 1.2.6 FISH HABITAT | | |
| 2.6 Charrying and Nymagny Habitat | | |
| .2.6. Spawning and Nursery Habitat | | |
| Cable 5. Area Factors for Low Marsh, High Marsh, and Swan | p Communities. | |
| No. of ha of Fish Habitat | Area Factor | |
| < 0.5 ha | 0.1 | |
| 0.5- 4.9 | 0.2 | |
| 5.0- 9.9 | 0.4 | |
| 10.0- 14.9 | 0.6 | |
| 15.0 -19.9 | 0.8 | |
| 20.0+ ha | 1.0 | |
| Step 1: | | |
| Fish habitat is not present within the wetland (Score = | 0) | |
| X Fish habitat is present within the wetland (Go to Step | 2) | |
| Step 2: Choose only one option | | |
| | | |
| Significance of the spawning and nursery habitation (Go to Step 3) | t within the wetland is known | |
| 2) X Significance of the spawning and nursery habitathem known (Go through Steps 4, 5, 6 and 7) | at within the wetland is not | |
| | | |
| | | |
| | | |
| | | |
| | | |

| No | orthern Onta | ario Wetland Evaluation | | Febru | ary 2012 | | | |
|---|---|---|------------|-------|----------|--|--|--|
| Step 3: Select the highest appropriate category below attach documentation: | | | | | | | | |
| 1) | | Significant in Site Region | 100 points | | | | | |
| 2) | | Significant in Site District | 50 | | | | | |
| 3) | | Locally Significant Habitat (5.0+ ha) | 25 | | | | | |
| 4) | | Locally Significant Habitat (<5.0 ha) | 15 | | | | | |
| | Score for Spawning and Nursery Habitat (maximum score 100 points) | | | | | | | |
| Step | <u>4:</u> Pro | oceed to Steps 4 to 7 <u>only</u> if Step 3 was <u>not</u> answ | vered. | | | | | |
| (Low | (Low Marsh: marsh area from the existing water line out to the outer boundary of the wetland) | | | | | | | |
| 2 | X Low marsh not present (Continue to Step 5) Low marsh present (Score as follows) | | | | | | | |

Scoring for Presence of Key Vegetation Groups

Scoring is based on the one most clearly dominant plant species of the dominant form in each Low Marsh vegetation community. Check the appropriate Vegetation Group (see Appendix 16) for each Low Marsh community. Sum the areas of the communities assigned to each Vegetation Group and multiply by the appropriate size factor from Table 5.

| Vegetation | Vegetation | Present | Total | Area | Score | Final |
|--------------|-----------------------------|---------------|-------|----------|-------|----------|
| Group Number | Group Name | as a | Area | Factor | | Score |
| | | Dominant | (ha) | | | (area |
| | | Form | | (see | | factor |
| | | (check) | | Table 5) | | x score) |
| | | | | | | |
| 1 | Tallgrass | | | | 6 pts | 0.0 |
| 2 | Shortgrass-Sedge | | | | 11 | 0.0 |
| 3 | Cattail-Bulrush-Burreed | | | | 5 | 0.0 |
| 4 | Arrowhead-Pickerelweed | | | | 5 | 0.0 |
| 5 | Duckweed | | | | 2 | 0.0 |
| 6 | Smartweed-Waterwillow | | | | 6 | 0.0 |
| 7 | Waterlily-Lotus | | | | 11 | 0.0 |
| 8 | Waterweed-Watercress | | | | 9 | 0.0 |
| 9 | Ribbongrass | | | | 10 | 0.0 |
| 10 | Coontail-Naiad-Watermilfoil | | | | 13 | 0.0 |
| 11 | Narrowleaf Pondweed | | | | 5 | 0.0 |
| 12 | Broadleaf Pondweed | | | | 8 | 0.0 |
| | Total Score (max | imum 75 point | s) | | | 0.0 |

| 1 | V | orthern | Ont | tario | WA | tland | 1 Eva | hrati | ۸r |
|---|---|---------|-----|-------|-----|-------|--------|-------|-------|
| 1 | N | OHIDEHI | CHI | ano | VVC | панс | i izva | шаш | . , , |

February 2012

Step 5: (**High Marsh**: area from the water line to the inland boundary of marsh wetland type. This is essentially what is commonly referred to as a wet meadow, in that there is insufficient standing water to provide fisheries habitat except during flood or high water conditions.)

High marsh not present (Continue to Step 6)

X High marsh present (Score as follows)

Scoring for Presence of Key Vegetation Groups

Scoring is based on the one most clearly dominant plant species of the dominant form in each High 1Marsh vegetation community. Check the appropriate Vegetation Group for each High Marsh community. Sum the areas of the communities assigned to each Vegetation Group and multiply by the appropriate size factor from Table 5.

| Vegetation | Vegetation | Present | Total | Area | Score | Final | | |
|---------------------------------|-------------------------|----------|-------|----------|-------|----------|--|--|
| Group Number | Group Name | as a | Area | Factor | | Score | | |
| | | Dominant | (ha) | (see | | (area | | |
| | | Form | | Table 5) | | factor | | |
| | | (check) | | | | x score) | | |
| 1 | Tallgrass | | | | 6 pts | 0.0 | | |
| 2 | Shortgrass-Sedge | X | 10.84 | 0.6 | 11 | 6.6 | | |
| 3 | Cattail-Bulrush-Burreed | | | | 5 | 0.0 | | |
| 4 | Arrowhead-Pickerelweed | | | | 5 | 0.0 | | |
| Total Score (maximum 25 points) | | | | | | | | |

Step 6: (**Swamp**: Swamp communities containing fish habitat, either seasonally or permanently. Determine the total area of seasonally flooded swamps and permanently flooded swamps containing fish habitat.)

X Swamp containing fish habitat not present (Continue to Step 7)
Swamp containing fish habitat present (Score as follows)

| Swamp containing fish Habitat | Present (check) | Total area (ha) | Area Factor (see Table 5) | | TOTAL SCORE (factor x score) |
|----------------------------------|--------------------|--------------------|------------------------------|----|---------------------------------|
| Seasonally flooded | | | | 10 | 0.0 |
| Permanently flooded | | | | 10 | 0.0 |
| SCO | RE (maximu | m 20 points) |) | | 0.0 |

| Northern Ontario Wetland Evaluation | February 2012 |
|---|--------------------------------------|
| Step 7: Calculation of final score | |
| Score for Spawning and Nursery Habitat (Low Marsh) (maximum 75) | = 0.0 |
| Score for Spawning and Nursery Habitat (High Marsh) (maximum 25) | = 6.6 |
| Score for Swamp Containing Fish Habitat (maximum 20) | = 0.0 |
| | imum score 100 points) = 6.6 |
| 4.2.6.2 Migration and Staging Habitat | • / |
| Step 1: | |
| 1) Staging or Migration Habitat is not present in the wetland (| (Score = 0) |
| 2) Staging or Migration Habitat is present in the wetland significant significant states and significant significant states are significant significant states. | ificance of the habitat is known (Go |
| to Step 2) 3) X Staging or Migration Habitat is present in the wetland significant (Go to Step 3) | ificance of the habitat is not known |
| NOTE: Only <u>one</u> of Step 2 <u>or</u> Step 3 is to be scored. | |
| Step 2: Select the highest appropriate category below, attach docum | |
| 1) Significant in Site Region | Score 25 points |
| 2) Significant in Site District | 15 |
| 3) Locally Significant | 10 |
| Fish staging and/or migration habitat present,but not as above | 5 |
| Score for Fish Migration and Staging Habitat (ma | eximum score 25 points) |
| Step 3: Select the highest appropriate category below based on pre (does not have to be dominant). Note name of river for 2) and 3). | sence of the designated site type |
| Wetland is riverine at rivermouth or lacustrine at rivermouth | Score h 25 points |
| 2) Wetland is riverine, within 0.75 km of rivermouth | 15 |
| Wetland is lacustrine, within 0.75 km of rivermouth | 10 |
| Fish staging and/or migration habitat present, but not as above | 5 |
| Score for Staging and Migration Habitat (ma | aximum score 25 points) 25 |
| | <u>-</u> |
| 22 | |

Northern Ontario Wetland Evaluation

February 2012

4.3 ECOSYSTEM AGE

(Fractional Area = area of wetland type/total area of wetland)

| | Area | | | Scoring |
|----------------------------------|------|------------|------|---------|
| Bog | | X | 25 = | 0.0 |
| Fen, treed to open on deep soils | | • | | |
| floating mats or marl | | X | 20 = | 0.0 |
| Fen, on limestone rock | | X | 5 = | 0.0 |
| Swamp | 0.87 | X | 3 = | 2.6 |
| Marsh | 0.13 | X | 0 = | 0.0 |
| | | Sub Total: | | 2.6 |

Fractional

Ecosystem Age Score (maximum 25 points)

2.6

4.4 GREAT LAKES COASTAL WETLANDS

Score for **coastal** (see text for definition) wetlands only

Choose one only

 wetland < 10 ha</td>
 = 0 points

 wetland 10- 50 ha
 = 25

 wetland 51 -lOO ha
 = 50

 wetland > 100 ha
 = 75

Great Lakes Coastal Wetlands Score (maximum 75 points)

| Northern Ontario Wetland Evaluation, Data and Sco | oring Reco | ord | February 2012 |
|--|------------|---|---------------|
| 5.0 EXTRA INFORMATION | | | |
| | | | |
| 5.1 PURPLE LOOSESTRIFE | | | |
| X Absent/Not seen | | | |
| Present Present | (a) | One location in wetland Two to many locations | <u> </u> |
| | (b) | Abundance code (1 < 20 plants (2 20-99 plants (3 100-999 plants (4 >1000 plants | |
| 5.2 SEASONALLY FLOODED AREAS | | | |
| Indicate length of seasonal flooding | | | |
| Check one or more | | | |
| Ephemeral Temporal Seasonal Semi-permanent No seasonal flooding | | (less than 2 weeks) (2 weeks to 1 month) (1 to 3 months) (>3 months) | <u>X</u> |
| 5.3 SPECIES OF SPECIAL SIGNIFICANCE | | | |
| 5.3.1 Osprey | | | |
| Present and nesting (attach map showing nest site) Known to have nested in last 5 yr Feeding area for osprey Not as above | | <u>X</u> | |
| 5.3.2 Common Loon | | | |
| Nesting in wetland (attach map showing nest site) Feeding at edge of wetland Observed or heard on lake or river adjoining the wetland Not as above | | X | |
| | | | |
| | | | |
| | | | |
| | 34 | | |

| Northern Ontario Wetland Evaluation, Data and Scoring Record | February 2012 |
|---|---------------------------------------|
| INVESTIGATORS | <u>AFFILIATION</u> |
| David Stephenson | Natural Resource Solutions Inc. |
| Charlotte Moore | Natural Resource Solutions Inc. |
| Jessica Grealey | Natural Resource Solutions Inc. |
| Katharina Walton | Natural Resource Solutions Inc. |
| Megan Pope | Natural Resource Solutions Inc. |
| Tara Brenton | Natural Resource Solutions Inc. |
| Tutu Divinon | Trada Resource Solutions Inc. |
| DATES WETLAND VISITED June 21 and 22, 2011 | |
| | |
| DATE THIS EVALUATION COMPLETED: | February 22, 2012 |
| ESTIMATED TIME DEVOTED TO COMPLETING THE FIELD 50 hours | SURVEY IN "PERSON HOURS" |
| | |
| WEATHER CONDITIONS | |
| i) at time of field work June 21 morning: 13°C, 70-90% cloud co June 21 evening: 15°C, 5-15% cloud cover, wind – Beaufort scale June 22: 10-24°C, 10-100% cloud cover, wind – Beaufort sc | 2-4 |
| ii) summer conditions in general spring: wet, cool; summer: ho | rt, dry |
| OTHER POTENTIALLY USEFUL INFORMATION: | , , , , , , , , , , , , , , , , , , , |
| Surveys completed by Natural Resource Solutions Inc.: | |
| vegetation, breeding birds, nocturnal birds, anuran call surveys | |
| vegetation, orecame on as, noctarnal on as, anaran can surveys | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| CHECKLIST OF PLANT AND ANIMAL SPECIES RECORDED IN T | HE WETLAND: |
| Attach a list of all flora and fauna observed in the wetland. | |
| *Indicate if voucher specimens or photos have been obtained, where local | ated, etc. |
| | |
| | |
| 35 | |

| North | nern Ontario Wetland Evaluation | | February 2012 |
|----------------------------------|---|-------------------------------------|-------------------------------|
| | WETLAND I | EVALUATION SCORING RECORD | |
| WETLAND | NAME | Abitibi-Martin's Meadow-Empire Wetl | and Complex |
| | <u>1.0 E</u> | BIOLOGICAL COMPONENT | |
| 1.1 | PRODUCTIVITY | | |
| 1.1.2 | Growing Degree-Days/Soils Wetland Type Site Type | | 11 9 2 |
| | | Total for Productivity | 22 |
| 1.2 | BIODIVERSITY | | |
| 1.2.2 1.2.3 1.2.4 1.2.5 | Number of Wetland Types Vegetation Communities (maxixmu Diversity of Surrounding Habitat (r Proximinty to Other Wetlands Interspersion Open Water Type | | 13 13 7 8 24 8 |
| 1.3 | Sub Total for Biodiversity <u>SIZE</u> (Biological Component) | Total for Biodiversity 73 | 73 37 |
| <u>TOT</u> 4 | AL FOR BIOLOGICAL COMPONE | NT (not to exceed 250) | 132 |
| | | | |

| Northern Ontario Welland Evaluation | February 2012 |
|--|---------------------------|
| 2.0 SOCIAL COMPONENT | |
| 2.1 ECONOMICALLY VALUABLE PRODUCTS | |
| 2.1.1 Wood Products 2.1.2 Lowbush Cranberry 2.1.3 Wild Rice 2.1.4 Commercial Fish 2.1.6 Furbearers | 14 0 10 12 12 |
| Total for Economically Valuable Products | 48 |
| 2.2 RECREATIONAL ACTIVITIES (maximum 80) | 16 |
| 2.3 LANDSCAPE AESTHETICS | |
| 2.3.1 Distinctness 2.3.2 Absence of Human Disturbance | <u>0</u> 4 |
| Total for Landscape Aesthetics | 4 |
| 2.4 EDUCATION AND PUBLIC AWARENESS | |
| 2.4.1 Educational Uses 2.4.2 Facilities and Programs 2.4.3 Research and Studies (maximum 12) | 0 0 0 |
| Total for Education and Public Awareness | 0 |
| 2.5 PROXIMITY TO AREAS OF HUMAN SETTLEMENT | 16 |
| 2.6 OWNERSH1P Subtotal for Social Component 2.7 SIZE (Social Component) | 19 |
| 2.8 ABORIGINAL AND CULTURAL VALUES (maximum 30) | 0 |
| TOTAL FOR SOCIAL COMPONENT (not to exceed 250) | 107 |
| | |
| | |
| | |
| | |
| | |

| Northern Ontario Wetland Evaluation, Score Summary | Februa | ary 2012 |
|--|---------------|----------|
| 3.0 HYDROLOGICAL COMPONENT | | |
| 3.1 <u>FLOOD ATTENUATION</u> | | 90 |
| 3.2 <u>GROUNDWATER RECHARGE</u> | | |
| 3.2.1 Site Type 3.2.2 Soils | 17 7 | |
| Total for Groundwater Recharge | | 24 |
| 3.3 <u>WATER QUALITY IMPROVEMENT</u> | | |
| 3.3.1 Watershed Improvement Factor3.3.2 Adjacent and Watershed Land Use3.3.3 Vegetation Form | 30 10 8 | |
| Total for Water Quality Improvement | | 48 |
| 3.4 <u>CARBON SINK</u> | | 9 |
| 3.5 SHORELINE EROSION CONTROL | | 8 |
| 3.6 GROUNDWATER DISCHARGE | | 26 |
| TOTAL FOR HYDROLOGICAL COMPONENT (not to exceed 250) | | 205 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

| Northern Ontario Wetland Evaluation, Score Summary | February 2012 |
|--|------------------------------------|
| 4.0 SPECIAL FEATURES | |
| 4.1 <u>RARITY</u> | |
| 4.1.1 Wetlands | 40 |
| 4.1.2 Species 4.1.2.1 Endangered or Threatened Species Breeding 4.1.2.2 Traditional Use by Endangered or Threatened Species 4.1.2.3 Provincially Significant Animals 4.1.2.4 Provincially Significant Plants 4.1.2.5 Regionally Significant Species 4.1.2.6 Locally Significant Species 4.1.2.7 Species of Special Status | 0 0 0 0 50 0 |
| Total for Species Rarity | 50 |
| 4.2 <u>SIGNIFICANT FEATURES OR HABITAT</u> | |
| 4.2.1 Colonial Waterbirds 4.2.2 Winter Cover for Wildlife 4.2.3 Waterfowl Staging and Moulting 4.2.4 Waterfowl Breeding 4.2.5 Migratory Passerine, Shorebird or Raptor Stopover 4.2.6 Ungulate Habitat 4.2.7 Fish Habitat | 0 0 0 10 0 25 32 |
| Total for Significant Features and | Habitat <u>67</u> |
| 4.3 ECOSYSTEM AGE | 3 |
| 4.4 <u>GREAT LAKES COASTAL WETLANDS</u> | 0 |
| TOTAL FOR SPECIAL FEATURES (maximum 250 | 159 |
| | |


| Nort | hern Ontario Wetland Evaluation, Score Summa | ury | February 2012 |
|-------------|--|----------------------------|---------------|
| | SUMMARY OF EV | ALUATION RESULT | |
| Wetland | Abitibi-Martin's Mead | low-Empire Wetland Complex | |
| TOTAL FO | OR 1.0 BIOLOGICAL COMPONENT | | 132 |
| TOTAL FO | OR 2.0 SOCIAL COMPONENT | | 107 |
| TOTAL FO | DR 3.0 HYDROLOGICAL COMPONENT | | 205 |
| TOTAL FO | DR 4.0 SPECIAL FEATURES COMPONENT | | 159 |
| | | WETLAND TOTAL | 603 |
| INVESTIG | <u>SATORS</u> | | |
| | David Stephenson | | |
| | Charlotte Moore | | |
| | Jessica Grealey | | |
| | Katharina Walton | | |
| | Megan Pope | | |
| AFFILIAT | Tara Brenton | | |
| AFFILIAT | Natural Resource Solutions Inc. | | |
| <u>DATE</u> | February 22, 2012 | | |
| DITTE | 1 coldary 22, 2012 | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

| Species Observed | | Vegetation survey | Amphibian survey | Breeding bird survey | Nocturnal bird survey |
|------------------------------|------------------------------|----------------------|---------------------|-------------------------|--------------------------|
| Amphibians | | , , | | | |
| Mink frog | Rana septentrionalis | X | | | |
| Spring peeper | Pseudacris crucifer crucifer | | Х | | |
| Wood frog | Rana sylvatica | (Repo | ted by I | latch) | |
| | | | | | |
| Birds | | | | | |
| Alder flycatcher | Empidonax alnorum | Х | | Χ | |
| American crow | Corvus brachyrhynchos | X | | Χ | |
| American goldfinch | Carduelis tristis | X | | Χ | |
| American kestrel | Falco sparverius | Х | | | |
| American redstart | Setophaga ruticilla | | | Х | |
| American robin | Turdus migratorius | X | Х | Х | Х |
| Black and white warbler | Mniotilta varia | X | | Х | |
| Black-capped chickadee | Poecile atricapillus | Х | | Х | |
| Black-throated green warbler | Dendroica virens | Х | | Х | |
| Black-throated blue warbler | Denrioca caerulenscens | Х | | Х | |
| Blue jay | Cyanocitta cristata | | | Х | |
| Chestnut-sided warbler | Dendrioca pensylvanica | | | Х | |
| Common loon | Gavia immer | X | | Х | |
| Common yellowthroat | Geothlypis trichas | Х | | Х | |
| Eastern phoebe | Sayornis phoebe | Х | | Χ | |
| Gray catbird | Dumetella carolinensis | | | Χ | |
| Hermit thrush | Catharus guttatus | | Х | Х | Х |
| Mourning warbler | Oporornis philadelphia | | | Х | |
| Northern cardinal | Cardinalis cardinalis | | | Х | |
| Nothern harrier | Circus cyaneus | Х | | | |
| Ovenbird | Seiurus aurocapillus | | | Х | |
| Red-eyed vireo | Vireo olivaceus | Х | | Х | |
| Red-winged blackbird | Agelaius phoeniceus | Х | | Х | |
| Ring-billed gull | Larus delawarensis | X | | | |
| Sandhill crane | Grus canadensis | Х | Х | Х | Х |
| Savannah sparrow | Passerculus sandwichensis | | | Х | |
| Scarlet tanager | Piranga olivacea | | | Х | |
| Sharp-shinned hawk | Accipiter striatus | Х | | | |
| Song sparrow | Melospiza melodia | | | Х | |
| Tennesee warbler | Vermivora peregrina | | | Х | |
| Tree swallow | Tachycineta bicolor | Х | | | |
| Veery | Catharus fuscescens | X | Х | Х | Х |
| White-breasted nuthatch | Sitta carolinensis | Х | | | |
| White-throated sparrow | Zonotrichia albicollis | X | Х | Х | Х |
| Yellow rumped warbler | Dendroica coronata | | | Х | |
| Yellow warbler | Dendroica petechia | Х | | Х | |
| | · | | | | |
| | | | | | |

| Species Observed | | Vegetation survey | Amphibian survey | Breeding bird survey | Nocturnal bird survey |
|----------------------------|--------------------------------------|----------------------|---------------------|-------------------------|--------------------------|
| Butterflies | | | | | |
| Canadian tiger swallowtail | Papilio canadensis | Х | | | |
| Common ringlet | Coenonympha tullia | X | | | |
| Juvenal's duskywing | Erynnis juvenalis | Х | | | |
| Northern crescent | Phyciodes pascoensis | X | | | |
| White admiral | Limenitis arthemis arthemis | X | | | |
| Wild indigo duskywing | Erynnis Baptisiae | Х | | | |
| Dragonflies and Darners | | | | | |
| Ebony jewelwing | Calopteryx maculata | Х | | | |
| Mammals | | | | | |
| Beaver | Castor canadensis | Х | | | |
| Groundhog | Marmota monax | X | | | |
| Moose | Alces alces | X | 1 | Х | 1 |
| Red fox | Vulpes vulpes | | | X | Х |
| Red squirrel | Tamiasciurus hudsonicus | Х | | | |
| White-tailed deer | Odocoileus virginianus | X | | Х | |
| willte-tailed deel | Odoconeds virginiarius | | | _^_ | |
| Vegetation | | | | | |
| Alder-leaved buckthorn | Rhamnus alnifolia | Х | | | |
| Aquatic sedge | Carex aquatilsis | Х | | | |
| Awl-fruited sedge | Carex stipata | Х | | | |
| Balsam fir | Abies balsamea | Х | | | |
| Balsam poplar | Populus balsamifera ssp. balsamifera | X | | | |
| Bebb's willow | Salix bebbiana | X | | | |
| Bird's-foot trefoil | Lotus corniculatus | X | | | |
| Black spruce | Picea mariana | X | | | |
| Black willow | Salix nigra | Х | | | |
| Blue bells | Campanula rotundifolia | X | | | |
| Blue flag iris | Iris versicolor | X | | | |
| Bluebead-lily | Clintonia borealis | X | | | |
| Bottlebrush sedge | Carex hystericina | Х | | | |
| Bracken fern | Pteridium aquilinum var. latiusculum | X | | | |
| Bristly black currant | Ribes lacustre | X | | | |
| Bull thistle | Cirsium vulgare | X | | | |
| Bunchberry | Cornus canadensis | X | | | |
| Bush honeysuckle | Diervilla Ionicera | X | | | |
| Canada blue-joint | Calamagrostis canadensis | X | | | |
| Canada mayflower | Maianthemum canadense | X | | | |
| Canada soapberry | Shepherdia canadensis | Х | | | |
| Choke cherry | Prunus virginiana ssp. virginiana | Х | | | |
| Club moss sp. | Lycopodiaceae sp. | Х | | | |
| Common cattail | Typha latifolia | Х | | | |
| Common dandelion | Taraxacum officinale | Х | | | |
| Common hairgrass | Deschampsia flexuosa | Х | | | |
| Cow parsnip | Heracleum maximum | Х | | | |
| Cow vetch | Vicia cracca | Х | | | |
| Curly dock | Rumex crispus | Х | | | |

| ecies Observed | | Vegetation survey | Amphibian survey | Breeding bird survey | Nocturnal |
|--------------------------------------|---|----------------------|---------------------|-------------------------|--|
| Dark-green bulrush | Scirpus atrovirens | X | | | |
| Dwarf raspberry | Rubus pubescens | Х | | | |
| Early meadowrue | Thalictrum dioicum | Х | | | |
| European moutain-ash | Sorbus aucuparia | Х | | | |
| Field horsetail | Equisetum arvense | Х | | | |
| Fireweed | Chamerion angustifolium spp. angustifolium | Х | | | |
| Fowl meadow grass | Glyceria striata | X | | | |
| Fox sedge | Carex vulpinoidea | Х | | | |
| Fragrant bedstraw | Galium triflorum | Х | | | |
| Grasses | Poa spp. | Х | | | |
| Greater duckweed | Spirodela polyrhiza | Х | | | |
| Hairy Solomon's seal | Polygonatum biflorum | Х | | | |
| High bush cranberry | Viburnum trilobum | Х | | | |
| Kentucky bluegrass | Poa saltuensis ssp. languida | Х | | | |
| Labrador-tea | Ledum groenlandicum | Х | | | |
| Lady fern | Athyrium filix-femina | Х | | | |
| Lettuce sp. | Lactuca sp. | Х | | | |
| Long-leaved aster | Symphyotrichum robynsianum | Х | | | |
| Low bush blueberry | Vaccinium angustifolium | Х | | | |
| Marsh cinquefoil | Comarum palustre | Х | | | |
| Marsh St. John's-wort | Triadenum virginicum | Х | | | |
| Marsh-marigold | Caltha palustris | Х | | | |
| Moss sp. | | Х | | | |
| New England aster | Symphyotrichum novae-angliae | X | | | t |
| Nodding trillium | Trillium cernuum | Х | | | |
| Northern beech fern | Phegopteris connectilis | X | | | t |
| Ostrich fern | Matteuccia struthiopteris var. pensylvanica | X | | | t |
| Pale jewelweed | Impatiens pallida | X | | | t |
| Prickly rose | Rosa acicularis ssp. sayi | X | | | t |
| Red currant | Ribes rubrum | X | | | H |
| Red maple | Acer rubrum | X | | | |
| Red raspberry | Rubus idaeus ssp. idaeus | X | | | H |
| Red-berried elder | Sambucus racemosa ssp. pubens | X | | | |
| Red-osier dogwood | Cornus stolonifera | X | | | |
| Reed canary grass | Phalaris arundinacea | X | | | H |
| Rough-leaved goldenrod | Solidago patula | X | | | |
| Sarsaparilla | Aralia elata | X | | | ┢ |
| Sedge sp. | Carex sp. | X | | | |
| Serviceberry | Amelanchier humilis | X | | | |
| Showy mountain ash | Sorbus decora | X | | | _ |
| Small-fruited Bulrush | Scirpus microcarpus | X | | | _ |
| Smooth scouring-rush | Equisetum laevigatum | X | | | ₩ |
| Speckled alder | Alnus incana spp. rugosa | X | | | \vdash |
| Spinulose wood fern | Dryopteris carthusiana | X | | | \vdash |
| Spotted touch-me-not | Impatiens capensis | X | 1 | | \vdash |
| Star-flower | Trientalis borealis ssp. borealis | X | | | ⊬ |
| Stinging nettle | Urtica dioica | X | | | ₩ |
| | | X | | | |
| Swamp fly honeysuckle Tall buttercup | Lonicera oblongifolia Ranunculus acris | X | } | | <u> </u> |

| Species Observed | | | | Breeding bird survey | Nocturnal bird survey |
|-----------------------|-------------------------------|---|---------------------|-------------------------|--------------------------|
| Tall meadow-rue | Thalictrum pubescens | X | Amphibian survey | | |
| Tamarack | Larix laricina | Х | | | |
| Trembling aspen | Populus tremuloides | Х | | | |
| Tufted loosestrife | Lysimachia thyrsiflora | Х | | | |
| Tufted vetch | Vicia cracca | Х | | | |
| White birch | Betula papyrifera | Х | | | |
| White spruce | Picea glauca | Х | | | |
| Wild carrot | Daucus carota | Х | | | |
| Wild mint | Mentha arvensis ssp. borealis | X | | | |
| Wild strawberry | Fragaria virginiana | X | | | |
| Willow species | Salix species | Х | | | |
| Wood horsetail | Equisetum sylvaticum | Х | | | |
| Woodland strawberry | Fragaria vesca ssp. americana | Х | | | |
| Yellow lady's slipper | Cypridedium calceolus | Х | | | |







Amphibian Data Form Project No. 1247 UTM: Observer: JEO, Station Name: Abitibi 1 Date: June 21 Visit #: A Start time: 20:08 % Cloud cover: Air Temp: Wind speed: Water Water Temp: pH: Precipitation Description:

| CA | LL LEVEL CODES | Beaufort Wind Scale | | | |
|--|--|---------------------|---------|---|--|
| 1 | Calls can be counted; not simultaneous | 0 Calm | 0-2 | Smoke rises vertically | |
| 2 | Some simultaneous calls; distinguishable | 1 Light air | 3-5 | Smoke drifts, but wind vanes do not | |
| 3 | Calls not distinguishable individually overlapping | 2 Slight breeze | 6 – 11 | Wind felt on face, leaves rustle | |
| Enter as: Call code (# of individuals) e.g. 1 (2) | | 3 Gentle breeze | 12 – 19 | Leaves & small twigs in constant motion; light flags extended | |
| | | 4 Mod. breeze | 20 – 30 | Wind raises dust and loose paper; small branches move | |
| | | 5 Fresh breeze | 31 – 39 | Small trees in leaf begin to sway | |
| | | 6 Strong breeze | 40 - 50 | Large branches in motion; inconvenience felt when walking | |

100m

50m

225 Labrador Drive, Waterloo, Ontario, N2K 4M8 Tel: (519) 725-2227 Fax: (519) 725-2575 Web: www.nrsi.on.ca



| Project: Coch | Amphibian | Data Form | _Project No. | 247 |
|-----------------------|--------------------------|--------------|------------------|----------------|
| Observer: JEG | Station Name: / Visit #: | | Date:3 | une 21. |
| Wind speed: 4 | % Cloud cover: | Air Temp: | Water Temp: / | Water pH: / |
| Precipitation Descrip | tion: none | | | |
| Remarks: | | | | |
| | (| directionO_° | | |
| | | 5 F | PE/(3) | |
| | | | (3) |) |
| | | | | |
| | | | | |
| | | | | |
| | | | | 1 |

50m 100m

| CALL LEVEL CODES | | Beaufor | Beaufort Wind Scale | | |
|--|--|--------------------|---------------------|--|--|
| 1 | Calls can be counted; not simultaneous | 0 Calm | 0-2 | Smoke rises vertically | |
| 2 | Some simultaneous calls; distinguishable | 1 Light air | 3-5 | Smoke drifts, but wind vanes do not | |
| 3 | Calls not distinguishable individually overlapping | 2 Slight breeze | 6 – 11 | Wind felt on face, leaves rustle | |
| Enter as: Call code (# of individuals) e.g. 1 (2) | | 3 Gentie breeze | 12 – 19 | Leaves & small twigs in constant motion; light flags extended | |
| | , | 4 Mod breeze | 20 – 30 | Wind raises dust and loose paper; small branches move | |
| | | 5 Fresh breeze | 31 – 39 | Small trees in leaf begin to sway | |
| | | 6 Strong breeze | 40 - 50 | Large branches in motion; inconvenience felt when walking against wind | |

225 Labrador Drive, Waterloo, Ontario, N2K 4M8 Tel: (519) 725-2227 Fax: (519) 725-2575 Web: www.nrsi.on.ca



| | Amphibian | Data F | orm | |
|------------------------------------|-----------------|------------|------------------------|----------------------|
| Project: Cock | rane | | Project No. | [24] |
| Observer: JEG | Station Name: A | | Dar Start tim | te:Junea 1e:20.29 |
| Wind speed: | % Cloud cover: | Air Tem | | 9°c pH:/ |
| Precipitation Descri | ption: | | | |
| Remarks: | | | | |
| | | direction_ | O . | |
| | | 5P | PE/CZ) | |
| | | | | |
| 7 | - | | | |
| 1/ | | | | |
| | | SPPE | 50 | Om 100m |
| CALL LEVEL CODES | Beaufor | rt Wind Sc | | |
| Calls can be counted; simultaneous | not 0 Calm | 0-2 | Śmoke rises vertically | |
| 2 Some simultaneous ca | alls; 1 Light | 3-5 | Smoke drifts, but wind | vanes do not |

| CALL LEVEL CODES | | Beaufort Wind Scale | | | |
|--|--|---------------------|---------|--|--|
| 1 | Calls can be counted; not simultaneous | 0 Calm | 0-2 | Smoke rises vertically | |
| 2 | Some simultaneous calls; distinguishable | 1 Light air | 3-5 | Smoke drifts, but wind vanes do not | |
| 3 | Calls not distinguishable individually overlapping | 2 Slight breeze | 6 – 11 | Wind felt on face, leaves rustle | |
| Enter as: Call code (# of individuals) e.g. 1 (2) | | 3 Gentle breeze | 12 – 19 | Leaves & small twigs in constant motion; light flags extended | |
| | | 4 Mod breeze | 20 – 30 | Wind raises dust and loose paper; small branches move | |
| | | 5 Fresh breeze | 31 – 39 | Small trees in leaf begin to sway | |
| | | 6 Strong breeze | 40 - 50 | Large branches in motion; inconvenience felt when walking against wind | |

225 Labrador Drive, Waterloo, Ontario, N2K 4M8 Tel: (519) 725-2227 Fax: (519) 725-2575 Web: www.nrsi.on.ca